


511,736
18 OCT 2004(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number
WO 03/103772 A1

- (51) International Patent Classification⁷: A61P 27/06, (74) Agent: OGILVY RENAULT; Attn: Murphy, Kevin, P., Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).
- (21) International Application Number: PCT/CA03/00838
- (22) International Filing Date: 2 June 2003 (02.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/386,499 6 June 2002 (06.06.2002) US
60/460,134 3 April 2003 (03.04.2003) US
- (71) Applicant (for all designated States except US):
MERCK FROSST CANADA & CO. [CA/CA]; 16711
Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BILLOT, Xavier [FR/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA). YOUNG, Robert, N. [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA). HAN, Yongxin [CA/CA]; 16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 1,5-DISTRIBUTED PYRROLID-2-ONE DERIVATIVES FOR USE AS EP₄ RECEPTOR AGONISTS IN THE TREATMENT OF EYE DISEASES SUCH AS GLAUCOMA

(57) Abstract: This invention relates to potent selective agonists of the EP₄ subtype of prostaglandin E₂ receptors of formula (I), their use or a formulation thereof in the treatment of glaucoma and other conditions which are related to elevated intraocular pressure in the eye of the patient.

BEST AVAILABLE COPY

WO 03/103772 A1

TITLE OF THE INVENTION

EP₄ RECEPTOR AGONIST, COMPOSITIONS AND METHODS THEREOF

This case claims the benefit of provisional application USSN 60/386,499, filed June 6, 2002 and USSN 60/460,134, filed April 3, 2003.

5

BACKGROUND OF THE INVENTION

Glaucoma is a degenerative disease of the eye wherein the intraocular pressure is too high to permit normal eye function. As a result, damage may occur to the optic nerve head and result in irreversible loss of visual function. If untreated, glaucoma may eventually lead to blindness. Ocular hypertension, i.e., the condition of elevated intraocular pressure without optic nerve head damage or characteristic glaucomatous visual field defects, is now believed by the majority of ophthalmologists to represent merely the earliest phase in the onset of glaucoma.

Many of the drugs formerly used to treat glaucoma proved unsatisfactory. Early methods of treating glaucoma employed pilocarpine and produced undesirable local effects that made this drug, though valuable, unsatisfactory as a first line drug. More recently, clinicians have noted that many β -adrenergic antagonists are effective in reducing intraocular pressure. While many of these agents are effective for this purpose, there exist some patients with whom this treatment is not effective or not sufficiently effective. Many of these agents also have other characteristics, e.g., membrane stabilizing activity, that become more apparent with increased doses and render them unacceptable for chronic ocular use and can also cause cardiovascular effects.

Agents referred to as carbonic anhydrase inhibitors decrease the formation of aqueous humor by inhibiting the enzyme carbonic anhydrase. While such carbonic anhydrase inhibitors are now used to treat elevated intraocular pressure by systemic and topical routes, current therapies using these agents, particularly those using systemic routes are still not without undesirable effects. Topically effective carbonic anhydrase inhibitors are disclosed in U.S. Patent Nos. 4,386,098; 4,416,890; 4,426,388; 4,668,697; 4,863,922; 4,797,413; 5,378,703, 5,240,923 and 5,153,192.

Prostaglandins and prostaglandin derivatives are also known to lower intraocular pressure. There are several prostaglandin types, including the A, B, C, D, E, F, G, I and J- Series (EP 0561073 A1). U.S. Patent 4,883,819 to Bito describes the use and synthesis of PGAs, PGBs and PGCs in reducing intraocular pressure. U.S. Patent 4,824,857 to Goh et al. describes the use and synthesis of PGD₂ and

derivatives thereof in lowering intraocular pressure including derivatives wherein C-10 is replaced with nitrogen. U.S. Patent 5,001,153 to Ueno et al. describes the use and synthesis of 13,14-dihydro-15-keto prostaglandins and prostaglandin derivatives to lower intraocular pressure. U.S. Patent 4,599,353 describes the use of eicosanoids and eicosanoid derivatives including prostaglandins and prostaglandin inhibitors in lowering intraocular pressure. See also WO 00/38667, WO 99/32441, WO 99/02165, WO 00/38663, WO 01/46140, EP 0855389, JP 2000-1472, US Patent No. 6,043,275 and WO 00/38690.

Prostaglandin and prostaglandin derivatives are known to lower intraocular pressure by increasing uveoscleral outflow. This is true for both the F type and A type of prostaglandins. This invention is particularly interested in those compounds that lower IOP via the uveoscleral outflow pathway and other mechanisms by which the E series prostaglandins (PGE₂) may facilitate IOP reduction. The four recognized subtypes of the EP receptor are believed to modulate the effect of lowering IOP (EP₁, EP₂, EP₃ and EP₄; *J. Lipid Mediators Cell Signaling*, Vol. 14, pages 83-87 (1996)). See also *J. Ocular Pharmacology*, Vol. 4, 1, pages 13-18 (1988); *J. Ocular Pharmacology and Therapeutics*, Vol. 11, 3, pages 447-454 (1995); *J. Lipid Mediators*, Vol. 6, pages 545-553 (1993); US Patent Nos. 5,698,598 and 5,462,968 and *Investigative Ophthalmology and Visual Science*, Vol. 31, 12, pages 2560-2567 (1990). Of particular interest to this invention are compounds, which are agonist of the EP₄ subtype receptor.

A problem with using prostaglandins or derivatives thereof to lower intraocular pressure is that these compounds often induce an initial increase in intraocular pressure, can change the color of eye pigmentation and cause proliferation of some tissues surrounding the eye.

As can be seen, there are several current therapies for treating glaucoma and elevated intraocular pressure, but the efficacy and the side effect profiles of these agents are not ideal. Therefore, there still exist the need for new and effective therapies with little or no side effects.

A variety of disorders in humans and other mammals involve or are associated with abnormal or excessive bone loss. Such disorders include, but are not limited to, osteoporosis, glucocorticoid induced osteoporosis, Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, bone fractures, rheumatoid arthritis, periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma. One of the most

common of these disorders is osteoporosis, which in its most frequent manifestation occurs in postmenopausal women. Osteoporosis is a systemic skeletal disease characterized by a low bone mass and microarchitectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture.

5 Osteoporotic fractures are a major cause of morbidity and mortality in the elderly population. As many as 50% of women and a third of men will experience an osteoporotic fracture. A large segment of the older population already has low bone density and a high risk of fractures. There is a significant need to both prevent and treat osteoporosis and other conditions associated with bone resorption. Because
10 osteoporosis, as well as other disorders associated with bone loss, are generally chronic conditions, it is believed that appropriate therapy will typically require chronic treatment.

Two different types of cells called osteoblasts and osteoclasts are involved in the bone formation and resorption processes, respectively. *See H. Fleisch,*
15 *Bisphosphonates In Bone Disease, From The Laboratory To The Patient*, 3rd Edition, Parthenon Publishing (1997), which is incorporated by reference herein in its entirety. Osteoblasts are cells that are located on the bone surface. These cells secrete an osseous organic matrix, which then calcifies. Substances such as fluoride, parathyroid hormone, and certain cytokines such as prostaglandins are known to provide a
20 stimulatory effect on osteoblast cells. However, an aim of current research is to develop therapeutic agents that will selectively increase or stimulate the bone formation activity of the osteoblasts.

Osteoclasts are usually large multinucleated cells that are situated either on the surface of the cortical or trabecular bone or within the cortical bone. The
25 osteoclasts resorb bone in a closed, sealed-off microenvironment located between the cell and the bone. The recruitment and activity of osteoclasts is known to be influenced by a series of cytokines and hormones. It is well known that bisphosphonates are selective inhibitors of osteoclastic bone resorption, making these compounds important therapeutic agents in the treatment or prevention of a variety of
30 systemic or localized bone disorders caused by or associated with abnormal bone resorption. However, despite the utility of bisphosphonates there remains the desire amongst researchers to develop additional therapeutic agents for inhibiting the bone resorption activity of osteoclasts.

Prostaglandins such as the PGE₂ series are known to stimulate bone
35 formation and increase bone mass in mammals, including man. It is believed that the

four different receptor subtypes, designated EP₁, EP₂, EP₃, and EP₄ are involved in mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts. The major prostaglandin receptor in bone is EP₄, which is believed to provide its effect by signaling via cyclic AMP.

5 In present invention it is further found that the formula I agonists of the EP₄ subtype receptor are useful for stimulating bone formation.


WO 02/24647, WO 02/42268, EP 1132086, EP 855389, EP 1114816, WO 01/46140 and WO 01/72268 disclose EP₄ agonists. However, they do not disclose the compounds of the instant invention.

10

SUMMARY OF THE INVENTION

This invention relates to potent selective agonists of the EP₄ subtype of prostaglandin E₂ receptors, their use or a formulation thereof in the treatment of glaucoma and other conditions that are related to elevated intraocular pressure in the
15 eye of a patient. Another aspect of this invention relates to the use of such compounds to provide a neuroprotective effect to the eye of mammalian species, particularly humans. This invention further relates to the use of the compounds of this invention for mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts.

20 More particularly, this invention relates to the use of EP₄ agonist having the structural formula I to treat ocular hypertension and/or glaucoma:





FORMULA I

25

or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug or mixture thereof, wherein,

X is (CH₂)_n, O or S;

Y represents $(C(R^b)_2)_n$, triple bond,  or  ;

R₁ represents hydroxy, CN, CHO, NHSO₂R₆, CONHSO₂R₆, CON(R₆)₂
 5 hydroxymethylketone, (CH₂)_pCO₂R₆, (CH₂)_nSO₃R₆, C₁₋₄ alkoxy, or (CH₂)_nC₅₋₁₀heterocyclyl, said heterocyclyl unsubstituted or substituted with 1 to 3 groups of R_a and optionally containing an acidic hydroxyl group, with the proviso that when X is a bond R₁ is not (CH₂)_pCO₂R₆, C₁₋₄ alkoxy, -(CH₂)_nNR₆R₇, CHO, NHSO₂R₆, CONHSO₂R₆, CON(R₆)₂, or hydroxymethylketone;

10

R₂ and R₃ independently represents hydrogen, or C₁₋₄ alkyl;

R₆ and R₇ independently represents hydrogen, or C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, (CH₂)_pC₆₋₁₀aryl, (CH₂)_pC₅₋₁₀heterocyclyl, CR₂R₃OC(O)OC₃₋₁₀cycloalkyl or
 15 CR₂R₃OC(O)O C₁₋₁₀alkyl;

Ar₂ independently represent (CH₂)_mC₆₋₁₀aryl, (CH₂)_mC₅₋₁₀heteroaryl, (CH₂)_mC₃₋₁₀ heterocycloalkyl, (CH₂)_mC₃₋₈ cycloalkyl said cycloalkyl, heterocycloalkyl, aryl or heteroaryl unsubstituted or substituted with 1-3 groups of R_a;

20

R_a represents C₁₋₆ alkoxy, C₁₋₆ alkyl, CF₃, nitro, amino, cyano, C₁₋₆ alkylamino, or halogen;

R^b independently represents H, halogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or

25

--- represents a double or single bond;

p represents 1-3;

30 n represents 0-4; and

m represents 0-8.

This and other aspects of the invention will be realized upon inspection of the invention as a whole.

DETAILED DESCRIPTION OF THE INVENTION

5 The invention is described herein in detail using the terms defined below unless otherwise specified.

 The term "therapeutically effective amount", as used herein, means that amount of the EP₄ receptor subtype agonist of formula I, or other actives of the present invention, that will elicit the desired therapeutic effect or response or provide
10 the desired benefit when administered in accordance with the desired treatment regimen. A preferred therapeutically effective amount relating to the treatment of abnormal bone resorption is a bone formation, stimulating amount. Likewise, a preferred therapeutically effective amount relating to the treatment of ocular
hypertension or glaucoma is an amount effective for reducing intraocular pressure
15 and/or treating ocular hypertension and/or glaucoma.

 "Pharmaceutically acceptable" as used herein, means generally suitable for administration to a mammal, including humans, from a toxicity or safety standpoint.

 The term "prodrug" refers to compounds which are drug
20 precursors which, following administration and absorption, release the claimed drug in vivo via some metabolic process. A non-limiting example of a prodrug of the compounds of this invention would be an acid of the pyrrolidinone group, where the acid functionality has a structure that makes it easily
hydrolyzed after administration to a patient. Exemplary prodrugs include acetic
25 acid derivatives that are non-narcotic, analgesics/non-steroidal, anti-inflammatory drugs having a free CH₂COOH group (which can optionally be in the form of a pharmaceutically acceptable salt, e.g. -CH₂COO-Na+), typically attached to a ring system, preferably to an aromatic or heteroaromatic ring system.

30 The term "alkyl" refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 10 carbon atoms unless otherwise defined. It may be straight, branched or cyclic. Preferred alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, t-butyl, cyclopentyl and cyclohexyl. When the alkyl group is said to be substituted with an alkyl group, this is used interchangeably with "branched alkyl
35 group".

Cycloalkyl is a specie of alkyl containing from 3 to 15 carbon atoms, without alternating or resonating double bonds between carbon atoms. It may contain from 1 to 4 rings, which are fused. Examples of cycloalkyl groups are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

5 Alkoxy refers to C₁-C₆ alkyl-O-, with the alkyl group optionally substituted as described herein. Examples of alkoxy groups are methoxy, ethoxy, propoxy, butoxy and isomeric groups thereof.

Halogen (halo) refers to chlorine, fluorine, iodine or bromine.

Aryl refers to aromatic rings e.g., phenyl, substituted phenyl and the like, as well as rings which are fused, e.g., naphthyl, phenanthrenyl and the like. An aryl group thus contains at least one ring having at least 6 atoms, with up to five such rings being present, containing up to 22 atoms therein, with alternating (resonating) double bonds between adjacent carbon atoms or suitable heteroatoms. The preferred aryl groups are phenyl, naphthyl and phenanthrenyl. Aryl groups may likewise be substituted as defined. Preferred substituted aryls include phenyl and naphthyl.

15 The term "heterocycloalkyl" refers to a cycloalkyl group (nonaromatic) having 3 to 10 carbon atoms in which one of the carbon atoms in the ring is replaced by a heteroatom selected from O, S or N, and in which up to three additional carbon atoms may be replaced by hetero atoms.

20 The term "cycloalkyl" refers to a cyclic alkyl group (nonaromatic) having 3 to 10 carbon atoms.

The term "heteroatom" means O, S or N, selected on an independent basis.



25 The term "heteroaryl" refers to a monocyclic aromatic hydrocarbon group having 5 or 6 ring atoms, or a bicyclic aromatic group having 8 to 10 atoms, containing at least one heteroatom, O, S or N, in which a carbon or nitrogen atom is the point of attachment, and in which one or two additional carbon atoms is optionally replaced by a heteroatom selected from O or S, and in which from 1 to 3 additional carbon atoms are optionally replaced by nitrogen heteroatoms, said heteroaryl group being optionally substituted as described herein. Examples of this type are pyrrole, pyridine, oxazole, thiazole, tetrazole, and oxazine. For purposes of this invention the tetrazole includes all tautomeric forms. Additional nitrogen atoms may be present together with the first nitrogen and oxygen or sulfur, giving, e.g., thiadiazole.

30 The term heterocyclyl or heterocyclic, as used herein, represents

a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from the group consisting of N, O, and S, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring. The term heterocycle or heterocyclic includes heteroaryl moieties. Examples of such heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnolinyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, 1,3-dioxolanyl, furyl, imidazolidinyl, imidazolyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholinyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, 2-oxopiperazinyl, 2-oxopiperdinyl, 2-oxopyrrolidinyl, piperidyl, piperazinyl, pyridyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxalinyl, tetrahydrofuryl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiazolyl, thiazolinyl, thienofuryl, thienothienyl, and thienyl. An embodiment of the examples of such heterocyclic elements include, but are not limited to, azepinyl, benzimidazolyl, benzisoxazolyl, benzofurazanyl, benzopyranyl, benzothiopyranyl, benzofuryl, benzothiazolyl, benzothienyl, benzoxazolyl, chromanyl, cinnolinyl, dihydrobenzofuryl, dihydrobenzothienyl, dihydrobenzothiopyranyl, dihydrobenzothiopyranyl sulfone, furyl, imidazolidinyl, imidazolyl, imidazolyl, indolinyl, indolyl, isochromanyl, isoindolinyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isothiazolidinyl, morpholinyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, 2-oxopiperazinyl, 2-oxopiperdinyl, 2-oxopyrrolidinyl, piperidyl, piperazinyl, pyridyl, 2-pyridinonyl, pyrazinyl, pyrazolidinyl, pyrazolyl, pyridazinyl, pyrimidinyl, pyrrolidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxalinyl, tetrahydrofuryl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiazolyl, thiazolinyl, thienofuryl, thienothienyl, thienyl and triazolyl.

For purposes of this invention, heterocyclyls containing acidic hydroxyl groups are those heterocyclyl groups that have an acidic hydroxy atom and

can have a pKa in the range of 3 to 7. Non-limiting examples of heterocyclcyls containing acidic hydroxyl groups are:



each R^c independently is H, fluorine,

cyano or C_{1-4} alkyl;

each R^d independently is H, C_{1-4} alkyl,

or a pharmaceutically acceptable cation;

each R^e independently is H, $-C(=O)-R^f$,

or $-SO_2R^e$, wherein R^f is C_{1-4} linear alkyl

or phenyl

The term "agonist" as used herein means EP4 subtype compounds of

- 5 formula I interact with the EP4 receptor to produce maximal, super maximal or submaximal effects compared to the natural agonist, PGE2. See Goodman and Gilman, The Pharmacological Basis of Therapeutics, 9th edition, 1996, chapter 2.



- 10 One embodiment of this invention is realized when compounds as described below are used to treat ocular hypertension and/or glaucoma. In particular, the compounds are those wherein R_1 is CN, $(CH_2)_n C_{5-10}$ heterocyclcyl, $(CH_2)_p CO_2R_6$ or $(CH_2)_n SO_3R_6$, said heterocyclcyl unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described. A subembodiment of this invention is realized when X is $(CH_2)_n$. Another subembodiment of this invention is when X is Sulfur. Still another sub-embodiment

of this invention is when X is oxygen. When R_1 is $(CH_2)_pCO_2R_6$, and X is sulfur, the sulfur is hexavalent.

Another embodiment of this invention is realized when Y is $(CH_2)_n$ and all other variables are as originally described.



5 Still another embodiment of this invention is realized when Y is $C(halo)_2$ and all other variables are as originally described.

Still another embodiment of this invention is realized when Y is a

double bond as described by  or  and all other variables are as originally described.

10 Still another embodiment of this invention is realized when Y is a triple bond and all other variables are as originally described.


Another embodiment of this invention is realized when R_1 is $(CH_2)_nC_{5-10}heterocyclyl$, said heterocyclyl unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described. A subembodiment of this invention is realized when X is $(CH_2)_n$, and Y is $(CH_2)_n$. Still another subembodiment of this invention is realized when X is $(CH_2)_n$, and Y is $C(halo)_2$. Yet another subembodiment of this invention is realized when X is a bond

and Y is a double bond as described by  or . Another subembodiment of this invention is realized when X is S.

20 Another embodiment of this invention is realized when R_1 is $(CH_2)_pCO_2R_6$, and all other variables are as originally described. A subembodiment of this invention is realized when X is $(CH_2)_n$, and Y is $(CH_2)_n$.

Another embodiment of this invention is realized when Ar_2 is $(CH_2)_mC_{6-10}aryl$, said aryl unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described.


25 Still another embodiment of this invention is realized when R_1 is a tetrazole unsubstituted or substituted with an R_a group and all other variables are as originally described. A subembodiment of this invention is realized when X is $(CH_2)_n$, and Y is $(CH_2)_n$, $C(halo)_2$ or a double bond as described by



Another subembodiment of this invention is realized when X is S.


Still another embodiment of this invention is realized when Ar₂ is a phenyl unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described.

Yet another embodiment of this invention is realized when R₁ is tetrazolyl and Ar₂ is phenyl, said tetrazolyl unsubstituted or substituted with a R_a group and phenyl is unsubstituted or substituted with 1-3 groups of R_a, and all other variables are as originally described. A subembodiment of this invention is realized when X is (CH₂)_n, and Y is (CH₂)_n, C(halo)₂ or a double bond as described by



Another subembodiment of this invention is realized when X is sulfur.

Still another embodiment of this invention is realized when a compound of formula II:



15

or a pharmaceutically acceptable salt, enantiomer, diastereomer, pro drug or mixture thereof is used, wherein X, Y, Ar₂ and n are as previously described.

A subembodiment of this invention is realized when X is (CH₂)_n, and

Y is (CH₂)_n, C(halo)₂ or a double bond as described by , and Ar₂ is phenyl. Another subembodiment of this invention is realized when n is 4.

A subembodiment of this invention is realized when X is S.

Compounds used in this invention are:

(5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)]butyl]pyrrolidin-2-one,

25

- 4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl cyanate,
 3-[4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]propanoic acid,
 5 [4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]methanesulfonic acid,
 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[1*H*-tetrazol-5-ylmethyl)butyl]pyrrolidin-2-one,
 [4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]acetic acid,
 10 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)thio]butyl]pyrrolidin-2-one,
 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(1*H*-tetrazol-5-ylthio)butyl]pyrrolidin-2-one,
 15 3-[4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]propanoic acid,
 [4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]methanesulfonic acid,
 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one,
 20 [4-[(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]acetic acid,
 (5*R*)-5[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-[6-(1*H*-tetrazol-5-yl)hexyl]pyrrolidin-2-one ,
 25 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
 7-[(2*S*)-2-[(3*R*)-4,4-difluoro-3-hydroxy-4-phenylbutyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 30 (5*Z*)-7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]hept-5-enoic acid,
 isopropyl (5*Z*)-7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]hept-5-enoate,

- 7-[(2*R*)-2-[(1*E*,3*R*)-4-(3-chlorophenyl)-4,4-difluoro-3-hydroxybut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4-(3-chlorophenyl)-4,4-difluoro-3-hydroxybut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 5 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-[3-(trifluoromethyl)phenyl]but-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-[3-(trifluoromethyl)phenyl]but-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
cyclopentyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 10 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-3-methyl-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-3-methyl-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 15 isobutyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate, and
cyclohexyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate, or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof.
- 20 Additional compounds are:
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)]butyl]pyrrolidin-2-one,
4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl cyanate,
- 25 3-[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]propanoic acid,
[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]methanesulfonic acid,
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[1*H*-tetrazol-5-ylmethyl]butyl]pyrrolidin-2-one,
- 30 4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]acetic acid,
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)thio]butyl]pyrrolidin-2-one,

(5*R*)-5-[(1*E*)- (3*R*) -hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(1*H*-tetrazol-5-ylthio)butyl]pyrrolidin-2-one,
3-[4-{(2*R*)-2-[(1*E*)- (3*R*) -hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoic acid,
5 [4-{(2*R*)-2-[(1*E*)- (3*R*) -hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]methanesulfonic acid,
(5*R*)-5-[(1*E*)- (3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one,
[4-{(2*R*)-2-[(1*E*)- (3*R*) -hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]acetic acid, or
10 (5*R*)-5-[(1*E*)-4,4-difluoro-(3*R*) -hydroxy-4-phenylbut-1-enyl]-1-[6-(1*H*-tetrazol-5-yl)hexyl]pyrrolidin-2-one, a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof.

Another embodiment of this invention is directed to a composition
15 containing an EP4 agonist of Formula I and optionally a pharmaceutically acceptable carrier.

Yet another embodiment of this invention is directed to a method for decreasing elevated intraocular pressure or treating glaucoma by administration, preferably topical or intra-cameral administration, of a composition containing an EP4
20 agonist of Formula I and optionally a pharmaceutically acceptable carrier. Use of the compounds of formula I for the manufacture of a medicament for treating elevated intraocular pressure or glaucoma or a combination thereof is also included in this invention.

This invention is further concerned with a process for making a
25 pharmaceutical composition comprising a compound of formula I.

This invention is further concerned with a process for making a pharmaceutical composition comprising a compound of formula I, and a pharmaceutically acceptable carrier.

The claimed compounds bind strongly and act on PGE2 receptor,
30 particularly on the EP4 subtype receptor and therefore are useful for preventing and/or treating glaucoma and ocular hypertension.

Dry eye is a common ocular surface disease afflicting millions of people. Although it appears that dry eye may result from a number of unrelated pathogenic causes, the common end result is the breakdown of the tear film, which
35 results in dehydration of the exposed outer surface of the eye. (Lemp, Report of the

Nation Eye Institute/Industry Workshop on Clinical Trials in Dry Eyes, The *CLAO* Journal, 21(4):221-231 (1995)). One cause for dry eye is the decreased mucin production by the conjunctival cells and/or corneal epithelial cells of mucin, which protects and lubricates the ocular surface (Gipson and Inatomi, Mucin genes expressed by ocular surface epithelium. Progress in Retinal and Eye Research, 16:81-98 (1997)). Functional EP4 receptors have been found in human conjunctival epithelial cells (see US Patent 6,344,477, incorporated by reference in its entirety) and it is appreciated that both human corneal epithelial cells (Progress in Retinal and Eye Research, 16:81-98(1997)) and conjunctival cells (Dartt et al. Localization of nerves adjacent to goblet cells in rat conjunctiva. Current Eye Research, 14:993-1000 (1995)) are capable of secreting mucins. Thus, the compounds of formula I are useful for treating dry eye.

Macular edema is swelling within the retina within the critically important central visual zone at the posterior pole of the eye. An accumulation of fluid within the retina tends to detach the neural elements from one another and from their local blood supply, creating a dormancy of visual function in the area. It is believed that EP4 agonist which lower IOP are useful for treating diseases of the macular such as macular edema or macular degeneration. Thus, another aspect of this invention is a method for treating macular edema or macular degeneration.

Glaucoma is characterized by progressive atrophy of the optic nerve and is frequently associated with elevated intraocular pressure (IOP). It is possible to treat glaucoma, however, without necessarily affecting IOP by using drugs that impart a neuroprotective effect. See Arch. Ophthalmol. Vol. 112, Jan 1994, pp. 37-44; Investigative Ophthalmol. & Visual Science, 32, 5, April 1991, pp. 1593-99. It is believed that EP4 agonist which lower IOP are useful for providing a neuroprotective effect. They are also believed to be effective for increasing retinal and optic nerve head blood velocity and increasing retinal and optic nerve oxygen by lowering IOP, which when coupled together benefits optic nerve health. As a result, this invention further relates to a method for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof by using an EP4 agonist of formula I.

The compounds produced in the present invention are readily combined with suitable and known pharmaceutically acceptable excipients to produce compositions which may be administered to mammals, including humans, to achieve effective IOP lowering. Thus, this invention is also concerned with a method of

treating ocular hypertension or glaucoma by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a β -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol, carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a Maxi-K channel blocker as disclosed in USSN 60/389,205, filed June 17, 2002 (Attorney Docket 21121PV), 60/389,222, filed June 17, 2002 (Attorney docket 21092PV), 60/458,981, filed March 27, 2003 (Attorney docket 21101PV4), 60/424790, filed November 8, 2002 (Attorney docket 21260PV), 60/424808, filed November 8, 2002 (Attorney docket 21281PV), 09/765716, filed January 17, 2001, 09/764738, filed January 17, 2001 and PCT publications WO 02/077168 and WO 02/02060863, all incorporated by reference in their entirety herein, a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as Iumigan and the compounds set forth in US Patent No. 5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; or an agonist of 5-HT₂ receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine.

This invention is also concerned with a method for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a β -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol, carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a Maxi-K channel blocker as disclosed in USSN 60/389,205, filed June 17, 2002 (Attorney Docket 21121PV), 60/389,222, filed June 17, 2002 (Attorney docket 21092PV), 60/458,981, filed March 27, 2003 (Attorney docket 21101PV4), 60/424790, filed November 8, 2002 (Attorney docket 21260PV), 60/424808, filed November 8, 2002 (Attorney docket 21281PV), 09/765716, filed January 17, 2001, 09/764738, filed January 17, 2001 and PCT

publications WO 02/077168 and WO 02/02060863, all incorporated by reference in their entirety herein, a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as lumigan and the compounds set forth in US Patent No. 5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; or an agonist of 5-HT₂ receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine. Use of the compounds of formula I for the manufacture of a medicament for increasing retinal and optic nerve head blood velocity, or increasing retinal and optic nerve oxygen tension or providing a neuroprotective effect or a combination thereof is also included in this invention.

This invention is further concerned with a method for treating macular edema or macular degeneration by administering to a patient in need thereof one of the compounds of formula I alone or in combination with a β -adrenergic blocking agent such as timolol, betaxolol, levobetaxolol, carteolol, levobunolol, a parasympathomimetic agent such as pilocarpine, a sympathomimetic agents such as epinephrine, iopidine, brimonidine, clonidine, para-aminoclonidine, a carbonic anhydrase inhibitor such as dorzolamide, acetazolamide, metazolamide or brinzolamide; a Maxi-K channel blocker as disclosed in USSN 60/389,205, filed June 17, 2002 (Attorney Docket 21121PV), 60/389,222, filed June 17, 2002 (Attorney docket 21092PV), 60/458,981, filed March 27, 2003 (Attorney docket 21101PV4), 60/424790, filed November 8, 2002 (Attorney docket 21260PV), 60/424808, filed November 8, 2002 (Attorney docket 21281PV), 09/765716, filed January 17, 2001, 09/764738, filed January 17, 2001 and PCT publications WO 02/077168 and WO 02/02060863, all incorporated by reference in their entirety herein, a prostaglandin such as latanoprost, travaprost, unoprostone, rescula, S1033 (compounds set forth in US Patent Nos. 5,889,052; 5,296,504; 5,422,368; and 5,151,444); a hypotensive lipid such as lumigan and the compounds set forth in US Patent No. 5,352,708; a neuroprotectant disclosed in US Patent No. 4,690,931, particularly eliprodil and R-eliprodil as set forth in WO 94/13275, including memantine; or an agonist of 5-HT₂ receptors as set forth in PCT/US00/31247, particularly 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate and 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-

ethylamine. Use of the compounds of formula I for the manufacture of a medicament for macular edema or macular degeneration is also included in this invention.

The EP4 agonist used in the instant invention can be administered in a therapeutically effective amount intravenously, subcutaneously, topically, 5 transdermally, parenterally or any other method known to those skilled in the art. Ophthalmic pharmaceutical compositions are preferably adapted for topical administration to the eye in the form of solutions, suspensions, ointments, creams or as a solid insert. Ophthalmic formulations of this compound may contain from 0.001 to 5% and especially 0.001 to 0.1% of medicament. Higher dosages as, for example, 10 up to about 10% or lower dosages can be employed provided the dose is effective in reducing intraocular pressure, treating glaucoma, increasing blood flow velocity or oxygen tension. For a single dose, from between 0.001 to 5.0 mg, preferably 0.005 to 2.0 mg, and especially 0.005 to 1.0 mg of the compound can be applied to the human eye.

15 The pharmaceutical preparation which contains the compound may be conveniently admixed with a non-toxic pharmaceutical organic carrier, or with a non-toxic pharmaceutical inorganic carrier. Typical of pharmaceutically acceptable carriers are, for example, water, mixtures of water and water-miscible solvents such as lower alkanols or aralkanols, vegetable oils, peanut oil, polyalkylene glycols, 20 petroleum based jelly, ethyl cellulose, ethyl oleate, carboxymethyl-cellulose, polyvinylpyrrolidone, isopropyl myristate and other conventionally employed acceptable carriers. The pharmaceutical preparation may also contain non-toxic auxiliary substances such as emulsifying, preserving, wetting agents, bodying agents and the like, as for example, polyethylene glycols 200, 300, 400 and 600, carbowaxes 25 1,000, 1,500, 4,000, 6,000 and 10,000, antibacterial components such as quaternary ammonium compounds, phenylmercuric salts known to have cold sterilizing properties and which are non-injurious in use, thimerosal, methyl and propyl paraben, benzyl alcohol, phenyl ethanol, buffering ingredients such as sodium borate, sodium acetates, gluconate buffers, and other conventional ingredients such as sorbitan 30 monolaurate, triethanolamine, oleate, polyoxyethylene sorbitan monopalmitate, dioctyl sodium sulfosuccinate, monothioglycerol, thiosorbitol, ethylenediamine tetracetic acid, and the like. Additionally, suitable ophthalmic vehicles can be used as carrier media for the present purpose including conventional phosphate buffer vehicle systems, isotonic boric acid vehicles, isotonic sodium chloride vehicles, isotonic 35 sodium borate vehicles and the like. The pharmaceutical preparation may also be in

the form of a microparticle formulation. The pharmaceutical preparation may also be in the form of a solid insert. For example, one may use a solid water soluble polymer as the carrier for the medicament. The polymer used to form the insert may be any water soluble non-toxic polymer, for example, cellulose derivatives such as methylcellulose, sodium carboxymethyl cellulose, (hydroxyloweralkyl cellulose), hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose; acrylates such as polyacrylic acid salts, ethylacrylates, polyactylamides; natural products such as gelatin, alginates, pectins, tragacanth, karaya, chondrus, agar, acacia; the starch derivatives such as starch acetate, hydroxymethyl starch ethers, hydroxypropyl starch, as well as other synthetic derivatives such as polyvinyl alcohol, polyvinyl pyrrolidone, polyvinyl methyl ether, polyethylene oxide, neutralized carbopol and xanthan gum, gellan gum, and mixtures of said polymer.

Suitable subjects for the administration of the formulation of the present invention include primates, man and other animals, particularly man and domesticated animals such as cats, rabbits and dogs.

The pharmaceutical preparation may contain non-toxic auxiliary substances such as antibacterial components which are non-injurious in use, for example, thimerosal, benzalkonium chloride, methyl and propyl paraben, benzyldodecinium bromide, benzyl alcohol, or phenylethanol; buffering ingredients such as sodium chloride, sodium borate, sodium acetate, sodium citrate, or gluconate buffers; and other conventional ingredients such as sorbitan monolaurate, triethanolamine, polyoxyethylene sorbitan monopalmitate, ethylenediamine tetraacetic acid, and the like.

The ophthalmic solution or suspension may be administered as often as necessary to maintain an acceptable IOP level in the eye. It is contemplated that administration to the mammalian eye will be from once up to three times daily.

For topical ocular administration the novel formulations of this invention may take the form of solutions, gels, ointments, suspensions or solid inserts, formulated so that a unit dosage comprises a therapeutically effective amount of the active component or some multiple thereof in the case of a combination therapy.

The compounds of the instant invention are also useful for mediating the bone modeling and remodeling processes of the osteoblasts and osteoclasts. See PCT US99/23757 filed October 12, 1999 and incorporated herein by reference in its entirety. The major prostaglandin receptor in bone is EP₄, which is believed to provide its effect by signaling via cyclic AMP. See Ikeda T, Miyaura C, Ichikawa A,

Narumiya S, Yoshiki S and Suda T 1995, *In situ localization of three subtypes (EP₁, EP₃ and EP₄) of prostaglandin E receptors in embryonic and newborn mice.*, *J Bone Miner Res* **10** (sup 1):S172, which is incorporated by reference herein in its entirety.

Use of the compounds of formula I for the manufacture of a medicament for
5 mediating the bone modeling and remodeling processes are also included in this invention

Thus, another object of the present invention is to provide methods for stimulating bone formation, i.e. osteogenesis, in a mammal comprising administering to a mammal in need thereof a therapeutically effective amount of an EP₄ receptor
10 subtype agonist of formula I.

Still another object of the present invention to provide methods for stimulating bone formation in a mammal in need thereof comprising administering to said mammal a therapeutically effective amount of an EP₄ receptor subtype agonist of formula I and a bisphosphonate active. Use of the compounds of formula I for the
15 manufacture of a medicament for stimulating bone formation is also included in this invention.

Yet another object of the present invention to provide pharmaceutical compositions comprising a therapeutically effective amount of an EP₄ receptor subtype agonist of formula I and a bisphosphonate active.

It is another object of the present invention to provide methods for
20 treating or reducing the risk of contracting a disease state or condition related to abnormal bone resorption in a mammal in need of such treatment or prevention, comprising administering to said mammal a therapeutically effective amount of an EP₄ receptor subtype agonist of formula I. Use of the compounds of formula I for the
25 manufacture of a medicament for treating or reducing the risk of contracting a disease state or condition related to abnormal bone resorption is also included in this invention.

The disease states or conditions related to abnormal bone resorption include, but are not limited to, osteoporosis, glucocorticoid induced osteoporosis,
30 Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, bone fractures, rheumatoid arthritis, periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma.

Within the method comprising administering a therapeutically effective amount of an EP₄ receptor subtype agonist of formula I and a bisphosphonate active,

both concurrent and sequential administration of the EP₄ receptor subtype agonist of formula I and the bisphosphonate active are deemed within the scope of the present invention. Generally, the formulations are prepared containing 5 or 10 mg of a bisphosphonate active, on a bisphosphonic acid active basis. With sequential
5 administration, the agonist and the bisphosphonate can be administered in either order. In a subclass of sequential administration the agonist and bisphosphonate are typically administered within the same 24 hour period. In yet a further subclass, the agonist and bisphosphonate are typically administered within about 4 hours of each other.

10 Nonlimiting examples of bisphosphonate actives useful herein include the following:

Alendronic acid, 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid;

Alendronate (also known as alendronate sodium or alendronate
15 monosodium trihydrate), 4-amino-1-hydroxybutylidene-1,1-bisphosphonic acid monosodium trihydrate;

Alendronic acid and alendronate are described in U.S. Patents 4,922,007, to Kieczkowski et al., issued May 1, 1990; 5,019,651, to Kieczkowski et al., issued May 28, 1991; 5,510,517, to Dauer et al., issued April
20 23, 1996; 5,648,491, to Dauer et al., issued July 15, 1997, all of which are incorporated by reference herein in their entirety;

Cycloheptylaminomethylene-1,1-bisphosphonic acid, YM 175, Yamanouchi (cimadronate), as described in U.S. Patent 4,970,335, to Isomura et al., issued November 13, 1990, which is incorporated by reference herein in its
25 entirety;

1,1-dichloromethylene-1,1-diphosphonic acid (clodronic acid), and the disodium salt (clodronate, Procter and Gamble), are described in Belgium Patent 672,205 (1966) and *J. Org. Chem* 32, 4111 (1967), both of which are incorporated by reference herein in their entirety;

30 1-hydroxy-3-(1-pyrrolidiny)-propylidene-1,1-bisphosphonic acid (EB-1053);

1-hydroxyethane-1,1-diphosphonic acid (etidronic acid);

1-hydroxy-3-(N-methyl-N-pentylamino)propylidene-1,1-bisphosphonic acid, also known as BM-210955, Boehringer-Mannheim

(ibandronate), is described in U.S. Patent No. 4,927,814, issued May 22, 1990, which is incorporated by reference herein in its entirety;

6-amino-1-hydroxyhexylidene-1,1-bisphosphonic acid
(neridronate);

5 3-(dimethylamino)-1-hydroxypropylidene-1,1-bisphosphonic acid
(olpadronate);

3-amino-1-hydroxypropylidene-1,1-bisphosphonic acid
(pamidronate);

[2-(2-pyridinyl)ethylidene]-1,1-bisphosphonic acid (piridronate) is
10 described in U.S. Patent No. 4,761,406, which is incorporated by reference in its
entirety;

1-hydroxy-2-(3-pyridinyl)-ethylidene-1,1-bisphosphonic acid
(risedronate);

(4-chlorophenyl)thiomethane-1,1-disphosphonic acid (tiludronate)
15 as described in U.S. Patent 4,876,248, to Breliere et al., October 24, 1989, which
is incorporated by reference herein in its entirety; and

1-hydroxy-2-(1H-imidazol-1-yl)ethylidene-1,1-bisphosphonic acid
(zolendronate).

A non-limiting class of bisphosphonate actives useful in the instant
20 invention are selected from the group consisting of alendronate, cimadronate,
clodronate, tiludronate, etidronate, ibandronate, neridronate, olpadronate,
risedronate, piridronate, pamidronate, zolendronate, pharmaceutically acceptable salts
thereof, and mixtures thereof.

A non-limiting subclass of the above-mentioned class in the instant
25 case is selected from the group consisting of alendronate, pharmaceutically acceptable
salts thereof, and mixtures thereof.

A non-limiting example of the subclass is alendronate monosodium
trihydrate.

In the present invention, as it relates to bone stimulation, the agonist is
30 typically administered for a sufficient period of time until the desired therapeutic
effect is achieved. The term "until the desired therapeutic effect is achieved", as used
herein, means that the therapeutic agent or agents are continuously administered,
according to the dosing schedule chosen, up to the time that the clinical or medical
effect sought for the disease or condition being mediated is observed by the clinician
35 or researcher. For methods of treatment of the present invention, the compounds are

continuously administered until the desired change in bone mass or structure is observed. In such instances, achieving an increase in bone mass or a replacement of abnormal bone structure with normal bone structure are the desired objectives. For methods of reducing the risk of a disease state or condition, the compounds are
5 continuously administered for as long as necessary to prevent the undesired condition. In such instances, maintenance of bone mass density is often the objective.

Nonlimiting examples of administration periods can range from about 2 weeks to the remaining lifespan of the mammal. For humans, administration periods can range from about 2 weeks to the remaining lifespan of the human,
10 preferably from about 2 weeks to about 20 years, more preferably from about 1 month to about 20 years, more preferably from about 6 months to about 10 years, and most preferably from about 1 year to about 10 years.


The instant compounds are also useful in combination with known agents useful for treating or preventing bone loss, bone fractures, osteoporosis,
15 glucocorticoid induced osteoporosis, Paget's disease, abnormally increased bone turnover, periodontal disease, tooth loss, osteoarthritis, rheumatoid arthritis, , periprosthetic osteolysis, osteogenesis imperfecta, metastatic bone disease, hypercalcemia of malignancy, and multiple myeloma. Combinations of the presently disclosed compounds with other agents useful in treating or preventing osteoporosis
20 or other bone disorders are within the scope of the invention. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful based on the particular characteristics of the drugs and the disease involved. Such agents include the following: an organic bisphosphonate; a cathepsin K inhibitor; an estrogen or an estrogen receptor modulator; an androgen receptor modulator; an
25 inhibitor of osteoclast proton ATPase; an inhibitor of HMG-CoA reductase; an integrin receptor antagonist; an osteoblast anabolic agent, such as PTH; calcitonin; Vitamin D or a synthetic Vitamin D analogue; and the pharmaceutically acceptable salts and mixtures thereof. A preferred combination is a compound of the present invention and an organic bisphosphonate. Another preferred combination is a
30 compound of the present invention and an estrogen receptor modulator. Another preferred combination is a compound of the present invention and an estrogen. Another preferred combination is a compound of the present invention and an androgen receptor modulator. Another preferred combination is a compound of the present invention and an osteoblast anabolic agent.

Regarding treatment of abnormal bone resorption and ocular disorders, the formula I agonists generally have an EC₅₀ value from about 0.001 nM to about 100 microM, although agonists with activities outside this range can be useful depending upon the dosage and route of administration. In a subclass of the present invention, the agonists have an EC₅₀ value of from about 0.01 microM to about 10 microM. In a further subclass of the present invention, the agonists have an EC₅₀ value of from about 0.1 microM to about 10 microM. EC₅₀ is a common measure of agonist activity well known to those of ordinary skill in the art and is defined as the concentration or dose of an agonist that is needed to produce half, i.e. 50%, of the maximal effect. See also, Goodman and Gilman's, *The Pharmacologic Basis of Therapeutics*, 9th edition, 1996, chapter 2, E. M. Ross, *Pharmacodynamics, Mechanisms of Drug Action and the Relationship Between Drug Concentration and Effect*, and PCT US99/23757, filed October 12, 1999, which are incorporated by reference herein in their entirety.

The herein examples illustrate but do not limit the claimed invention. Each of the claimed compounds are EP₄ agonists and are useful for a number of physiological ocular and bone disorders.

The compounds of this invention can be made, with some modification, in accordance with US Patent No. 6,043,275, EP0855389, USSN 60/337228 (Merck Docket No. MC052) and WO 01/46140, all of which are incorporated herein by reference in their entirety. The following non-limiting examples, given by way of illustration, is demonstrative of the present invention.


Scheme 1



The compounds embodied in this application may be synthesized in part by Scheme 1. Pyroglutamic acid is converted to the corresponding ester through the action of a dehydrating agent such as thienylchloride and an alcohol such as methanol. The ester can be reduced by a reducing agent such as sodium borohydrate to provide the corresponding alcohol. Protection of the alcohol with a suitable protecting group such as t-butyldimethylsilylchloride in the presence of a base such as imidazole provides these silyl-protected alcohol. Reacting the protected alcohol amide with a strong base such as sodium hydride and thence with an alkyl halide (RX) provides the N-alkylated product. Alternatively, a protecting group such as paramethoxybenzylbromide may be substituted for RX to provide a protected analog of the N-alkylated product. Removal of the silyl-protecting group with a reagent such as HF-pyridine and then subsequent oxidation of the free alcohol with an oxidizing agent such as 2-periodinane then provides the aldehyde 1.

An aryl ketoester such as 2 is reacted with a fluoronating agent such as DAST to provide the difluoro ester 3. Reaction with lithiomethylene dimethoxyphosphonate provides the ketophosphonate 4. Reaction of the aldehyde 1 with the ketophosphonate 4 in the presence of a strong base such as sodium hydride provides the olefin 5. Reduction of the ketone group with a reducing agent such as sodium borohydride provides the alcohol 6. The alcohol 6 may subsequently be protected with a suitable protecting group such as t-butyldimethylsilylchloride and thence, if desired, the protected amide (such as paramethoxybenzyl = R) it would be deprotected utilizing an oxidizing agent such as ceric ammonium nitrate. The free amide then be reacted with a strong base (sodium hydride) and then reacted with a suitably elaborated alkyl halide RX to provide the final products.

Scheme 2




5 An alternative scheme is shown in Scheme 2. Glutamic acid is treated with an aromatic aldehyde such as paramethoxybenzylaldehyde, in the presence of reducing agents such as sodium borohydride or sodium cyanoborohydride to provide the N-alkylated product 7. Cyclization and ester synthesis are accomplished using methanol and acidic catalysts to afford 8. Reduction of the ester to the alcohol 9 is accomplished using a reducing agent such as sodium borohydride and then oxidation

of the alcohol moiety with common oxidation reagents such as pyridine, sulfur trioxide, etc. provides the aldehyde 10. Reaction of the aldehyde 10 with difluoroketophosphonate 4 in the presence of a strong base such as LDA or sodium hydride allows the formation of enone 11. The protecting group R_1 can then be removed in an oxidation step using a reagent such as cerium ammonium nitrate to afford the lactam 12. The enone moiety is reduced to the allyl alcohol using a common reducing agent such as sodium borohydride and then protected in the usual manner by reaction with t-butyldimethylsilylchloride to afford a protected alcohol 13. The amide 13 can then be alkylated with a variety of groups R_3 by reaction first with a strong base such as sodium hydride in a polar non protic solvent in the presence of a phase-transfer catalyst such as tetrabutylammoniumiodide and then addition of electrophilic reagent R_3X where X is the halogen or a suitable leaving group to provide 14.


If R_3 is an aromatic group, the reaction conditions are that compound 13 is mixed with a base such as cesium carbonate in dioxane, the aromatic R_3X where X is halogen or triflate and a catalytic amount of palladium catalyst and the mixture is heated.

Scheme 3

20



Scheme 4



Preparation 1

Step A: 7-[2R-(tert-Butyl-dimethyl-silanyloxymethyl)-5-oxo-pyrrolidin-1-yl]-heptanenitrile.

Step B: 7-(2R-Hydroxymethyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile.

A solution of tetrabutylammonium fluoride (1M in THF, 100.0 mL, 100.0 mmol) was slowly added to a solution of 7-[2R-(tert-butyl-dimethyl-silanyloxymethyl)-5-oxo-pyrrolidin-1-yl]-heptanenitrile (22.39 g, 66.13 mmol) in THF (400 mL) at 0°C. The reaction was warmed to room temperature and was stirred for 4 h. Saturated aqueous NaHCO₃ (250 mL) was added and the volatiles were removed in vacuo. The remaining aqueous solution was washed with CHCl₃ (4x200 mL). The combined organic solutions were dried (MgSO₄), filtered, and concentrated. Purification by medium pressure chromatography eluting with a solvent gradient (9:1 hexanes:EtOAc to 4:1 hexanes:EtOAc to 7:3 hexanes:EtOAc to 6:4 hexanes:EtOAc to 1:1 hexanes:EtOAc to EtOAc to 9:1 EtOAc:MeOH) provided

7-(2R-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile (14.922 g). ^1H NMR (CDCl_3) δ 3.78 (dd, 1 H), 3.71-3.58 (m, 3H), 3.00 (m, 1 H), 2.46 (m, 1 H), 2.36-2.27 (m, 3H), 2.08 (m, 1 H), 1.93 (m, 1 H), 1.77 (m, 1 H), 1.68-1.43 (m, 6H), 1.32 (m, 2H); MS 225.1 (M+1).

5

Step C:

7-(2R-formyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile.

- To a solution of 7-(2R-hydroxymethyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile (336 mg, 1.5 mmole) in CH_2Cl_2 (7 ml) was added portion-wise
- 10 Dess-Martin periodinane (636 mg, 1.5 mmole) and the reaction mixture was stirred 1h at rt. Solvent is removed in vacuo, and the residue is triturated with toluene, filtered on celite and solvent removed to give
- 7-(2R-formyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile as an oil.

Preparation 2

Dimethyl 3,3-difluoro-2-oxo-3-phenyl-propylphosphonate

To a solution of dimethyl methanephosphonate (1.139 g, 9.18 mmole) in 20 ml THF is added dropwise nBuLi (1.6 M in hexanes, 5.73 ml, 9.18 mmole) at -78°C. This solution is stirred 30 min at -78°C and then added to a solution of 2,2-difluorophenylacetic acid methyl ester (1.75 g, 8.74 mmole) at -78°C. The reaction mixture is allowed to reach rt and then acetic acid (1.5 ml) and 10 ml water are added. The aqueous phase is extracted three times with 30 ml AcOEt, the organic phases are then washed with water, brine, dried on Na₂SO₄ and the solvent is removed under reduced pressure. Purification of the residual oil by silica gel flash chromatography (3:7 Acetone : toluene) to give Dimethyl 3,3-difluoro-2-oxo-3-phenyl-propylphosphonate as an oil. ¹H NMR (CDCl₃) δ 7.65-7.40 (m, 5 H), 3.71-3.58 (m, 3H), 3.77 (s, 3 H), 3.74 (s, 3 H), 3.35 (d, *J* = 22 Hz, 2H); MS 279.1 (M+1).

Preparation 3

(5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-pyrrolidin-2-one

Step A: (5*R*)-(tert-butyl-dimethyl-silanyloxymethyl)-1-(4-chlorobutyl)pyrrolidin-2-one

To a solution of (5*R*)-(tert-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one (*Tetrahedron: Asymmetry*, **1996**, 7, 2113) (2.83 g, 12.34 mmol) in 60 ml DMF was added NaH (95%, 325.7 mg, 13.57 mmol) in one portion and the mixture was heated at 50°C for 30 min. Then 4-bromo-1-chlorobutane (2.96 g, 17.27 mmol) and a catalytic amount of nBu₄NI were added and the mixture was stirred at 50°C for 1 h. The reaction was cooled to room temperature and water (100 ml) was added. The aqueous phase was extracted with AcOEt (4x200ml), the organic phases were washed with water (200 ml), brine (100 ml), dried on MgSO₄, filtered and the solvent was removed under reduced pressure. The residual oil was purified by flash column-chromatography on silica gel (eluent AcOEt 1: Hexanes 1) to provide (5*R*)-(tert-butyl-dimethyl-silanyloxymethyl)-1-(4-chlorobutyl)pyrrolidin-2-one as an oil. ¹H NMR (CDCl₃) 3.71-3.53 (m, 6H), 3.05 (m, 1H), 2.46-2.24 (m, 2H), 2.05 (m, 1H), 1.84-1.61 (m, 4H), 0.85 (s, 9H), 0.03 (s, 6H); MS 320.2-322.2 (M+1).

Step B: (5R)-1-(4-chlorobutyl)-5-(hydroxymethyl)pyrrolidin-2-one

To a solution of (5R)-(tert-butyl-dimethyl-silanyloxymethyl)-1-(4-chlorobutyl)pyrrolidin-2-one (1.95 g, 6.11 mmol) in CH₂Cl₂ (25 ml) in a Teflon Erlenmeyer at 0°C was added dropwise HF-pyridine complex (1 ml), and the solution was allowed to reach room temperature, and was stirred for 1.5 h. Water (20 ml) and 1N HCl (1 ml) were added to the reaction mixture. The aqueous phase was extracted with CH₂Cl₂ (4x30ml), the organic phases was washed with brine (20 ml), dried on MgSO₄, filtered and the solvent was removed under reduced pressure. The residual oil was purified by flash column-chromatography on silica gel (eluent Acetone 1: Toluene 1) to provide (5R)-1-(4-chlorobutyl)-5-(hydroxymethyl)pyrrolidin-2-one as an oil. ¹H NMR (CDCl₃) 4.00 (s, 1H), 3.71-3.53 (m, 6H), 3.03 (m, 1H), 2.46-2.22 (m, 2H), 2.14-1.88 (m, 2H), 1.79-1.55 (m, 4H); MS 206.1-208.1 (M+1).

Step C: (2R)-1-(4-chlorobutyl)-5-oxopyrrolidine-2-carboxaldehyde

To a solution of (5R)-1-(4-chlorobutyl)-5-(hydroxymethyl)pyrrolidin-2-one (309.6 mg, 1.5 mmol) in CH₂Cl₂ (7 ml) was added Dess-Martin periodinane (638 mg, 1.5 mmol) portionwise over 40 min at room temperature. After 1 h, the solvent was removed under reduced pressure, and the residue triturated with Et₂O (3x5 ml), filtered on a celite plug, and the solvent removed. (2R)-1-(4-chlorobutyl)-5-oxopyrrolidine-2-carboxaldehyde was obtained as a colorless oil. ¹H NMR (CDCl₃) 9.58 (s, 1H), 4.18 (m, 1H), 3.65 (m, 1H), 3.53 (t, *J* = 8 Hz, 2H), 3.08 (m, 1H), 2.43 (m, 2H), 2.30(m, 1H), 2.08 (m, 1H), 1.78-1.56 (m, 4H).

Step D: (5R)-1-(4-chlorobutyl)-5-[(1E)-3-oxo-4-phenylbut-1-enyl]pyrrolidin-2-one

To a solution of (3-phenyl-2-oxo-propyl)-phosphonic acid dimethyl ester (938 mg, 4 mmol) in DME (20 ml) at 0°C was added portionwise NaH 95 % (100.8 mg, 4.2 mmol), and the mixture was stirred 20 min at 0°C. Then a solution of (2R)-1-(4-chlorobutyl)-5-oxopyrrolidine-2-carboxaldehyde in DME (5 ml) was added dropwise and the reaction mixture was allowed to reach room temperature, and stirred overnight. A half-saturated solution of NH₄Cl (10 ml) was added and the aqueous phase was extracted with AcOEt (4x60ml); the organic phases was washed with water (20 ml), brine (20 ml), dried on MgSO₄, filtered and the solvent was removed under reduced pressure. The residual oil was purified by flash column-chromatography on silica gel (eluent Acetone 2: Toluene 8) to provide (5R)-1-(4-chlorobutyl)-5-[(1E)-3-

oxo-4-phenylbut-1-enyl]pyrrolidin-2-one as an oil. ¹H NMR (CDCl₃) 7.35-7.20 (m, 5H), 6.64 (dd, J = 15.7 Hz, 8.2 Hz, 1H), 6.25 (d, J = 15.7 Hz, 1H), 4.17 (m, 1H), 3.85 (s, 2H), 3.55-3.50 (m, 3H), 2.77 (m, 1H), 2.43-2.17 (m, 3H), 1.81-1.75 (m, 1H), 1.70-1.51 (m, 4H).

5

Step E: (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]pyrrolidin-2-one

To a solution of (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-oxo-4-phenylbut-1-enyl]pyrrolidin-2-one (161 mg, 0.50 mmol) in MeOH (5 ml) at -20°C was added portionwise NaBH₄ (31 mg, 0.8 mmol). The mixture was stirred at -20°C for 1 h, and the solvent was removed under reduced pressure. The residue was dissolved in a mixture of water (5 ml) and 1N HCl (1 ml), the aqueous phase was extracted with AcOEt (3x15ml); the organic phases was washed with water (5 ml), brine (5 ml), dried on MgSO₄, filtered and the solvent was removed under reduced pressure. The residual oil was purified by flash column-chromatography on silica gel (eluent Acetone 4: Toluene 6) to provide both diastereoisomers of (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]pyrrolidin-2-one as an oil. ¹H NMR (CDCl₃) 7.36-7.22 (m, 5H), 5.78 (m, 1H), 5.51 (m, 1H), 4.44 (m, 1H), 4.07 (m, 1H), 3.59-3.45 (m, 3H), 2.95-2.77 (m, 3H), 2.44-2.19 (m, 3H), 2.43-2.17 (m, 3H), 1.70-1.55 (m, 5H).

20

Preparation 4

(5*R*)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-1-{4-[(triisopropylsilyl)thio]butyl}-pyrrolidin-2-one

To a solution of (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]pyrrolidin-2-one (273.9 mg, 0.852 mmol) in THF (5 ml) were added triisopropylsilylsulfide (324.4 mg, 1.70 mmol), a catalytic amount of nBu₄NI and portionwise NaH 95 % (30.7 mg, 1.28 mmol). The mixture was heated to 50°C for 1 h. The reaction was cooled to room temperature and water (2 ml) was added. The aqueous phase was extracted with AcOEt (4x10ml), the organic phases were washed with water (2 ml), brine (2 ml), dried on MgSO₄, filtered and the solvent was removed under reduced pressure. The residual oil was purified by flash column-chromatography on silica gel (eluent AcOEt 1: Hexanes 3) to provide both diastereoisomers of (5*R*)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-1-{4-[(triisopropylsilyl)thio]butyl}pyrrolidin-2-one as an oil. ¹H NMR (CDCl₃) 7.30-7.16 (m, 5H), 5.72 (m, 1H), 5.45 (m, 1H), 4.37 (m, 1H), 4.02 (m, 1H), 3.44 (m, 1H), 2.86-


35

2.79 (m, 3H), 2.51 (m, 2H), 2.35-2.12 (m, 3H), 1.94 (s, 1H), 1.64-1.50 (m, 5H), 1.2 (m, 3H), 1.05 (d, $J = 8.0$ Hz, 18H); MS 476.4 (M+1).

The difluoro component in the examples below, unless otherwise
 5 indicted, can be added to the following examples in accordance with schemes 1 and 2
 herein.

EXAMPLE 1


(5*R*)-5-[(1*E*)-3-hydroxy-difluoro-4-phenylbut-1-enyl]-1-{4-[(1-methyl-1*H*-tetrazol-
 10 5-yl)thio]butyl}pyrrolidin-2-one



To a solution of the (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-
 enyl]pyrrolidin-2-one (50.0 mg, 0.155 mmol) in DMF (0.5 ml) was added 5-
 15 mercapto-1-methyltetrazole sodium salt, and a catalytic amount of $n\text{Bu}_4\text{NI}$. The
 mixture is heated to 50°C overnight. The reaction is cooled to room temperature and
 water (5 ml) is added. The aqueous phase is extracted with AcOEt (4x10ml), the
 organic phases are washed with water (2 ml), brine (2 ml), dried on MgSO_4 , filtered
 and the solvent is removed under reduced pressure. The residual oil is purified by
 20 flash column-chromatography on silica gel (eluent AcOEt 2: Hexanes 3) to provide
 both diastereoisomers of (5*R*)-5-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-1-
 {4-[(1-methyl-1*H*-tetrazol-5-yl)thio]butyl}pyrrolidin-2-one.

EXAMPLE 2

4-[(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl
thiocyanate




5

To a solution of the (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]pyrrolidin-2-one (50.0 mg, 0.155 mmol) in DMF (1 ml) are added potassium
thiocyanate (150.7 mg, 1.55 mmol), and a catalytic amount of *n*Bu₄NI. The mixture is
10 heated to 50°C overnight. The reaction is cooled to room temperature and water (5
ml) is added. The aqueous phase is extracted with AcOEt (4x10ml), the organic
phases are washed with water (2 ml), brine (2 ml), dried on MgSO₄, filtered and the
solvent is removed under reduced pressure. The residual oil is purified by flash
column-chromatography on silica gel (eluent Acetone 4: Toluene 6) to provide both
15 diastereoisomers of 4-[(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-
oxopyrrolidin-1-yl]butyl thiocyanate.

EXAMPLE 3

20 3-[4-[(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-
yl]butyl]thio]propanoic acid



Step A : methyl 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoate

To a solution of the (5*R*)-1-(4-chlorobutyl)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]pyrrolidine-2-one (50.0 mg, 0.155 mmol) in DMF (0.8 ml) are added 3-mercaptopropanoic acid methyl ester (93.0 mg, 0.755 mmol), a catalytic amount of $n\text{Bu}_4\text{NI}$ and then dropwise 1M MeONa (0.62 ml, 0.62mmol). The mixture is heated to 80°C for 24h. The reaction is cooled to room temperature and water (6 ml) is added. The aqueous phase is extracted with AcOEt (4x10ml), the organic phases are washed with water (2 ml), brine (2 ml), dried on MgSO_4 , filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (eluent Acetone 4: Toluene 6) to provide both diastereoisomers of methyl 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoate.


Step B : 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoic acid

To a solution of methyl 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoate (19.6 mg, 0.0484 mmol) in MeOH/THF (1:1)(2 ml) is added a solution of 1N LiOH (0.051 ml, 0.051 mmol) at 0°C. The reaction mixture is stirred overnight at room temperature. 0.5N HCl (4 ml) is added, the aqueous phase is extracted with CH_2Cl_2 (4x10ml), the organic phases are washed with brine (2 ml), dried on MgSO_4 , filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (gradient CH_2Cl_2 : MeOH: AcOH (100:0:0) to (95:5:0.5)) to provide both diastereoisomers of 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]propanoic acid.

EXAMPLE 4

[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]methanesulfonic acid

5




To a solution of the (5*R*)-5-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-1-{4-[(triisopropylsilyl)thio]butyl}pyrrolidin-2-one (39.3 mg, 0.083 mmol) in THF (1 ml) are added sodium bromomethanesulfonate (32.6 mg, 0.165 mmol) and then dropwise 1M *n*Bu₄NF (0.25 ml, 0.25mmol). The mixture was heated to 50°C for 1h. The reaction is cooled to room temperature and 1N HCl (2 ml) is added. The aqueous phase is extracted with Et₂O (4x10ml), the organic phases are washed with 1N HCl (2 ml), brine (2 ml), dried on Na₂SO₄, filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (eluent CH₂Cl₂ 95: MeOH 5:AcOH 0.5) to provide both diastereoisomers of methyl [4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl)thio]methanesulfonic acid.

20

EXAMPLE 5

(5*R*)-5-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one



Step A : (5R)-5-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-1-[4-(methylthio)butyl]pyrrolidin-2-one


To a solution of the (5R)-5-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-1-[4-
 5 [(triisopropylsilyl)thio]butyl]pyrrolidin-2-one (46.2 mg, 0.097 mmol) in THF (1 ml)
 were added methyl iodide (17.6 mg, 0.126 mmol) and then dropwise 1M nBu₄NF
 (0.116 ml, 0.116mmol) at -78°C. The mixture is then stirred at room temperature for
 1h. NH₄Cl half saturated (2 ml) is added. The aqueous phase is extracted with AcOEt
 (5x8 ml), the organic phases are washed with brine (2 ml), dried on Na₂SO₄, filtered
 10 and the solvent is removed under reduced pressure. The residual oil is purified by
 flash column-chromatography on silica gel (eluent Acetone 4: Toluene 60) to provide
 (5R)-5-[(1E)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylthio)butyl]-
 pyrrolidin-2-one.

15 Step B: (5R)-5-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one

To a solution of (5R)-5-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-1-[4-
 (methylthio)butyl]pyrrolidin-2-one (31.2 mg, 0.093 mmol) in CH₂Cl₂ : MeOH :H₂O
 (7:2:1)(5 ml) was added portionwise Oxone[®] (172.9 mg, 0.281 mmol) at 0°C for 10
 20 min., and 4 h at room temperature. 5% solution of NaHSO₃ (2 ml) is added. The
 aqueous phase is extracted with CH₂Cl₂ (4x10 ml), the organic phases are washed
 with water (5 ml), brine (2 ml), dried on MgSO₄, filtered and the solvent is removed
 under reduced pressure. The residual oil is purified by flash column-chromatography
 on silica gel (eluent Acetone 7: Toluene 30) to provide (5R)-5-[(1E)-3-hydroxy-4-
 25 difluoro- 4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]pyrrolidin-2-one.

EXAMPLE 6

30 [4-{(2R)-2-[(1E)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butyl]thio]acetic acid



Step A: methyl (([4-((2R)-2-((tert-butyl(dimethyl)silyl)oxy)methyl)-5-oxopyrrolidin-1-yl]butyl)thio)acetate

To a solution of (5R)-(tert-butyl-dimethyl-silanyloxymethyl)-pyrrolidin-2-one (*Tetrahedron: Asymmetry*, 1996, 7, 2113) (1.5 g, 6.55 mmol) in 30 ml DMF is added NaH 95% (173.0 mg, 7.20 mmol) in one portion and the mixture is heated at 50°C for 30 min. Then 4-bromo-1-chlorobutane (1.347 g, 7.86 mmol) and a catalytic amount of $n\text{Bu}_4\text{NI}$ are added and the mixture is stirred at 50 °C for 1 h. The reaction is cooled to room temperature and methyl thioglycolate (1.39 g, 13.1 mmol), then dropwise addition of 4.9N MeONa (2.4 ml, 11.79 mmol). The mixture is stirred overnight at room temperature and water (150 ml) is added. The aqueous phase is extracted with AcOEt (4x150ml), the organic phases are washed with water (200 ml), brine (100 ml), dried on MgSO_4 , filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (eluent AcOEt 1: Hexanes 1) to provide methyl (([4-((2R)-2-((tert-butyl(dimethyl)silyl)oxy)methyl)-5-oxopyrrolidin-1-yl]butyl)thio)acetate.

Step B: methyl (([4-((2R)-2-(hydroxymethyl)-5-oxopyrrolidin-1-yl]butyl)thio)acetate

To a solution of methyl (([4-((2R)-2-((tert-butyl(dimethyl)silyl)oxy)methyl)-5-oxopyrrolidin-1-yl]butyl)thio)acetate (571 mg, 1.47 mmol) in CH_2Cl_2 (8 ml) in a Teflon Erlenmeyer at 0°C is added dropwise HF-pyridine complex (0.8 ml), and the solution is allowed to reach room temperature, and is stirred for 1.5 h.. Water (20 ml) and 1N HCl (1 ml) were added to the reaction mixture. The aqueous phase is extracted with CH_2Cl_2 (4x30ml), the organic phases is washed with brine (20 ml), dried on MgSO_4 , filtered and the solvent is removed under reduced pressure to provide methyl (([4-((2R)-2-(hydroxymethyl)-5-oxopyrrolidin-1-yl]butyl)thio)acetate.

Step C: Methyl ({4-[(2R)-2-formyl-5-oxopyrrolidin-1-yl]butyl}thio)acetate

To a solution of methyl ({4-[(2R)-2-(hydroxymethyl)-5-oxopyrrolidin-1-yl]butyl}thio)acetate (634.5 mg, 2.30 mmol) in CH₂Cl₂ (15 ml) is added Dess-Martin periodinane (975 mg, 1.5 mmol) portionwise over 40 min at room temperature. After 1 h, the solvent is removed under reduced pressure, and the residue triturated with Et₂O (3x5 ml), filtered on a Celite plug, and the solvent removed. Methyl ({4-[(2R)-2-formyl-5-oxopyrrolidin-1-yl]butyl}thio)acetate is obtained.

Step D: methyl [(4-{(5R)-2-oxo-5-[(1E)-3-oxo-4-phenylbut-1-enyl]pyrrolidin-1-yl}butyl)thio]acetate

To a solution of (3-phenyl-2-oxo-propyl)-phosphonic acid dimethyl ester (264 mg, 1.09 mmol) in DME (5 ml) at 0°C is added portionwise NaH 95 % (26 mg, 1.09 mmol), and the mixture is stirred 20 min at 0°C. Then a solution of methyl ({4-[(2R)-2-formyl-5-oxopyrrolidin-1-yl]butyl}thio) (270 mg, 0.99 mmol) in DME (2 ml) is added dropwise and the reaction mixture is allowed to reach room temperature, and stirred overnight. A half-saturated solution of NH₄Cl (5 ml) is added and the aqueous phase is extracted with AcOEt (4x10ml); the organic phases is washed with water (10 ml), brine (10 ml), dried on MgSO₄, filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (eluent AcOEt) to provide methyl [(4-{(5R)-2-oxo-5-[(1E)-3-oxo-4-phenylbut-1-enyl]pyrrolidin-1-yl}butyl)thio]acetate.

Step E: methyl [(4-{(2R)-2-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-5-pyrrolidin-1-yl}butyl)thio]acetate


To a solution of methyl [(4-{(5R)-2-oxo-5-[(1E)-3-oxo-4-phenylbut-1-enyl]pyrrolidin-1-yl}butyl)thio]acetate (295.6 mg, 0.75 mmol) in MeOH (5 ml) at -20°C was added portionwise NaBH₄ (27.6 mg, 1.2 mmol). The mixture is stirred at -20°C for 1 h, and the solvent is removed under reduced pressure. The residue is dissolved in a mixture of water (5 ml) and 1N HCl (1 ml), the aqueous phase is extracted with AcOEt (3x15ml); the organic phases is washed with water (5 ml), brine (5 ml), dried on MgSO₄, filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (eluent Acetone 4: Toluene 6) to provide both diastereoisomers of methyl [(4-{(2R)-2-[(1E)-3-hydroxy-4-phenylbut-1-enyl]-5-pyrrolidin-1-yl}butyl)thio]acetate.

Step F: [4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butylthio]acetic acid

- To a solution of methyl [(4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-phenylbut-1-enyl]-5-pyrrolidin-1-yl}butylthio)acetate (90.0 mg, 0.23 mmol) in MeOH/THF (1:2)(5 ml) is added a solution of 1*N* LiOH (0.46 ml, 0.46 mmol) at 0°C. The reaction mixture is stirred 4 h at room temperature. 1*N* HCl (3 ml) is added, the aqueous phase is extracted with CH₂Cl₂ (4x10ml), the organic phases are washed with brine (2 ml), dried on MgSO₄, filtered and the solvent is removed under reduced pressure. The residual oil is purified by flash column-chromatography on silica gel (gradient CH₂Cl₂: MeOH: AcOH (100:0:0) to (94:6:0.5)) to provide both diastereoisomers of [4-{(2*R*)-2-[(1*E*)-3-hydroxy-4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butylthio]-acetic acid.

EXAMPLE 7

- (5*R*)-5[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-[6-(1*H*-tetrazol-5-yl)hexyl]pyrrolidin-2-one



- Step A: 7-{2-Oxo-5*R*-[3-hydroxy-4,4-difluoro-4-phenyl)-but-1-enyl]-pyrrolidin-1-yl}-heptanenitrile.

- To a solution of dimethyl 3,3-difluoro-2-oxo-3-phenyl-propylphosphonate (405 mg, 1.46 mmole) in THF (5 ml) is added NaH (36.8 mg, 1.53 mmole) at 0°C, and the reaction mixture is stirred for 30 min at 0°C. A solution of 7-(2*R*-formyl-5-oxo-pyrrolidin-1-yl)-heptanenitrile (325 mg, 1.46 mmole) in THF (5 ml) is added to the above mixture. After 30 min at 0°C, the mixture is allowed to reach rt for 2h.

Addition of a saturated solution of ammonium chloride (5 ml), extraction with AcOEt (3x15 ml); the organic phases are washed with brine, dried on Na₂SO₄ and the solvent is removed. The residue is purified by a silica gel flash chromatography (70 : 30 AcOEt : Hexanes) to give 7-{2-Oxo-5*R*-[3-oxo-4,4-

difluoro-4- phenyl)-but-1-enyl]-pyrrolidin-1-yl}heptanenitrile as a oil. This oil is immediately dissolved in MeOH (5 ml), and $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (223 mg, 0.6 mmole) is added at -20°C . NaBH_4 (36.5 mg, 0.96 mmole) is added portionwise, and the mixture is allowed to reach rt. HCl 1N (1 ml) is added, solvent is removed in vacuo. The residue is dissolved in water (5 ml), extracted with AcOEt (3x10 ml) and the organic phases are washed with brine, dried on Na_2SO_4 and solvent removed. Purification by flash silica gel chromatography (30:70 Acetone : Toluene) to give 7-{2-Oxo-5R-[3-hydroxy-4,4-difluoro-4- phenyl)-but-1-enyl]-pyrrolidin-1-yl}-heptanenitrile as a oil. ^1H NMR (CDCl_3) δ 7.48-7.3 (m, 5 H), 5.65-5.61 (m, 1), 5.57-5.53 (m, 1h), 4.5 (m, 1 H), 3.48 (m, 1H), 3.35 (m, 1H), 2.68 (m, 1H), 2.32-2.20 (m, 5H), 2.10 (m, 1H), 1.55-1.15 (m, 8H); MS 377.3 (M+1).

Step B:

To 7-{2-Oxo-5R-[3-hydroxy-4,4-difluoro-4- phenyl)-but-1-enyl]-pyrrolidin-1-yl}-heptanenitrile (193.5 mg, 0.51 mmole) is added Tributylstannylazide (512 mg, 1.54 mmole) and the mixture is heated at 100°C for 8h. The reaction mixture is dissolved in AcOEt and worked-up with a mixture of 1:1 (1N HCl : 5% KF). The aqueous phase is extracted by AcOEt (3x10 ml), and the organic phases are washed with brine, dried on Na_2SO_4 and solvent removed. Purification by silica gel filtration, gradient (CH_2Cl_2 : MeOH : 0.1% HCOOH) from (100 : 0) to (94 : 6) to afford (5R)-5[(1E)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-[6-(1H-tetraazol-5-yl)hexyl]pyrrolidin-2-one as an oil. ^1H NMR (CDCl_3) δ 7.5-7.35 (m, 5H), 5.7 (m, 1H), 5.6 (m, 1H), 4.6 (m, 1H), 4.05 (m, H), 3.9 (m, 2H), 2.75 (m, 1H), 2.4 (m, 2H), 2.2 (m, 1H), 1.80-1.6 (m, 3H), 1.4-1.05 (m, 6H) ; MS 420.3 (M+1), 418.3 (M-1).

Examples 8-11 are prepared in accordance with Scheme 4.

EXAMPLE 8

isopropyl 7-{(2R)-2-[(1E,3R)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}heptanoate (8a)

Step A

(5R)-5-(hydroxymethyl)-1-(4-methoxybenzyl)pyrrolidin-2-one (1a)

The preparation of 1a was carried out according to the literature procedure (see: Tetrahedron 1994, 6221)

Step B

(5*R*)-5-[(1*E*)-4,4-difluoro-3-oxo-4-phenylbut-1-enyl]-1-(4-methoxybenzyl)pyrrolidin-2-one (2a)

5 Oxalyl chloride (2.75 mL) was added dropwise to a solution of dimethylsulfoxide (2.45 mL) in CH₂Cl₂ (60 ml) at -72 °C under a stream of N₂, and the mixture was stirred 20 min at that temperature. A solution of (5*R*)-5-(hydroxymethyl)-1-(4-methoxybenzyl)pyrrolidin-2-one (1a, 6.74 g) in CH₂Cl₂ (30 ml) was then added via a cannula, and the mixture was stirred at -72 °C for 20 min.
10 Triethylamine (8.45 ml) was then added and the mixture was stirred at -72 °C for an additional 30 min and then concentrated in vacuo (to remove most of the CH₂Cl₂). The residue was diluted with a mixture of ethyl acetate and diethyl ether and then filtered. The filtrate was concentrated to give the desired product. ¹H NMR (acetone-d₆) δ 9.50 (s, 1H), 7.20 (d, 2H), 6.90 (d, 2H), 4.80 (d, 1H), 4.14 (d, 1H), 4.06 (m, 1H),
15 3.79, (s, 3H), 2.20-2.45 (m, 3H), 2.12 (m, 1H).

 To a solution of dimethyl 3,3-difluoro-2-oxo-3-phenylpropylphosphonate (United States Patent 4,320,136 March. 16, 1982) (2.076 g) in THF (17 mL) at 0 °C was added potassium tert-butoxide (963 mg) and the mixture was stirred for an additional 1 hour at 0 °C. To the mixture was then added
20 (2*R*)-1-(4-methoxybenzyl)-5-oxopyrrolidine-2-carboxaldehyde in THF (10 mL) via cannula and the resultant mixture stirred at room temperature 2 hours and quenched with saturated NH₄Cl. The mixture was then extracted with ethyl acetate (3x) and the organic layer was washed with water, brine, dried over Mg₂SO₄, filtered and concentrated. The residue was purified by chromatography using 20% acetone/toluene
25 as the eluent to give the desired product 2a. ¹H NMR (400 MHz, CDCl₃): δ 7.60-7.45 (m, 5H), 7.00 (d, 2H), 6.78 (d, 2H), 6.45 (d, 1H), 4.90 (d, 1H), 4.10-4.00 (m, 1H), 3.80 (s, 3H), 3.70 (d, 1H), 3.35 (d, 1H), 2.55-2.40 (m, 2H), 2.25-2.15 (m, 1, 1H), 1.85-1.75 (m, 1H).

30 Step C

(5*R*)-5-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-(4-methoxybenzyl)pyrrolidin-2-one (3a)

 To a solution of (2a) (8.2 g, 21.2 mmol) in 80 mL CH₂Cl₂ was added (S)-CBS in toluene (10.6 mL, 10.6 mmol) and the mixture was cooled to -40 °C. A solution of catechol borane (6.8 mL, 63.8 mmol) in CH₂Cl₂ (20 mL) was added dropwise and
35 the solution was stirred at -40 °C for one hour, and allowed to warm up to -20 °C for

two hours. The reaction mixture was quenched at -20 °C with 1 N HCl and was stirred for 4 hours at room temperature. The phases were separated and the organic phase was sequentially washed with 1N HCl, H₂O, 1 N NaOH, brine and dried over Na₂SO₄, filtered and concentrated in vacuo. The crude was purified by flash chromatography using 40-50% ethyl acetate/hexanes to give the desired products as a mixture of diastereomers as a pale yellow oil. MS (M + 1) 388.2. The mixture of diastereomers was further separated by preparative HPLC (ChiralPak AD[®]). Eluting with 30% isopropyl alcohol in hexanes first gave isomer 4. Further elution afforded the major, more polar isomer **3a**.

Step D

(5*R*)-5-((1*E*,3*R*)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-4,4-difluoro-4-phenylbut-1-enyl)-1-(4-methoxybenzyl)pyrrolidin-2-one (**5a**)

To a solution **3a** (365 mg) in DMF (3 mL) at room temperature was added imidazole (139 mg) followed by TBSCl (220 mg). The mixture was stirred over the weekend and then quenched with water. The mixture was extracted with ether (3x) and washed with water, brine, dried over Na₂SO₄, filtered and concentrated in vacuo. Purified by column chromatography (50% ethyl acetate : hexane) afforded compound **5a**. ¹H NMR (400 MHz, CDCl₃): δ 7.55-7.40 (m, 5H), 7.15-7.10 (m, 2H), 6.85-6.80 (m, 2H), 5.65-5.55 (m, 2H), 4.95 (2d, 1H), 4.55-4.45 (m, 1H), 3.90-3.85 (m, 1H), 3.80 (s, 3H), 3.60 (2d, 1H), 2.50-2.30 (m, 2H), 2.20-2.05 (m, 1H), 1.70-1.55 (m, 1H), 0.85 (d, 9H), 0.00 (t, 6H).

Step E

(5*R*)-5-((1*E*,3*R*)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-4,4-difluoro-4-phenylbut-1-enyl)pyrrolidin-2-one (**6a**)

To a solution of **5a** (359 mg) in acetonitrile (20 mL) at 0 °C was added CAN (ceric(IV)ammonium nitrate) (2 g), water (2 mL) and the mixture was allowed to warm to room temperature for 4 hours. The mixture was extracted with ether (3x) and was washed with water, brine and dried over Na₂SO₄. Purification by column chromatography (50%-75%-100% ethyl acetate in hexane) afforded compound **5a**. ¹H NMR (400 MHz, CDCl₃): δ 7.50-7.40 (m, 5H), 5.70-5.65 (m, 2H), 4.50-4.42 (m, 1H), 4.20-4.13 (m, 1H), 2.37-2.30 (m, 3H), 1.80-1.70 (m, 1H), 0.87 (s, 9H), -0.05 (d, 6H).

Step F

isopropyl 7-{3-[(2*R*)-2-[(1*E*,3*R*)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate (7a)

To a solution of 5a (314 mg) in DMF (dimethyl formamide - 5 mL) was added NaH 60% (36.4 mg) and the mixture was stirred at room temperature for 1h until gas evolution was ceased. Iso-propyl 7-bromoheptanoate (415 mg) and a crystal of NaI was added. The mixture was heated to 90 °C for 6h. After cooling to room temperature, the mixture was quenched with saturated NH₄Cl and extracted with diethyl ether (3 x). The organic layer was washed with water, brine and dried over Na₂SO₄. The crude was purified by flash chromatography. Eluting with 50-60% ethyl acetate in hexanes gave the desired product 6a. ¹H NMR (400 MHz, acetone-d₆) δ 7.50 (m, 5H), 7.73, m, 2H), 4.96 (m, 1H), 4.70 (m, 1H), 4.18 (m, 1H), 3.44 (m, 1H), 2.74 (m, 1H), 2.30-2.18 (m, 5H), 1.70-1.20 (m, 9H), 1.20 (d, 6H), 0.87 (s, 9H), 0.05 (s, 3H) and -0.01 (s, 3H).

Step G

The title compound 8a

To a solution of 7a (350 mg) in THF (tetrahydrofuran - 5 mL) was added TBAF (1M in THF, 1.3 mL) and the mixture was stirred at room temperature for 30 min. The solution was then diluted with ethyl acetate and washed with water (4x) and brine. After drying over Na₂SO₄, the organic layer was filtered and concentrated in vacuo to give the desired title compound 8a. ¹H NMR (400 MHz, acetone-d₆) δ 7.56-7.47 (m, 5H), 5.79-5.65 (m, 2H), 5.05 (bs, 1H, OH), 4.95 (m, 1H), 4.66 (m, 1H), 4.12 (m, 1H), 3.30 (m, 1H), 2.70 (m, 1H), 2.30 -2.15 (m, 5H), 1.70-1.20 (m, 9H), 1.20 (d, 6H).

EXAMPLE 9

7-{(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}heptanoic acid (9a)

A mixture of compound 8a (94 mg) and LiOH (0.25 mL, 1M) in MeOH (1 mL) was stirred at room temperature overnight and concentrated in vacuo. The residue was co-evaporated with MeOH three times and the residue was washed with diethyl ether three times (to remove trace of unhydrolysed ester). After neutralizing with 1N HCl, the mixture was extracted with ethyl acetate (3x). The

organic layer was washed with water and brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated to give the title compound **9a**. MS (-ESI): m/z 394.3 (M-1).

EXAMPLE 10

5

Isopropyl 7-[(2*S*)-2-[(3*R*)-4,4-difluoro-3-hydroxy-4-phenylbutyl]-5-oxopyrrolidin-1-yl]heptanoate (**10a**)

To a solution of **8a** (77 mg) in degassed MeOH was added Pd/C (12 mg, 10%) and the mixture was purged with hydrogen three times and then stirred
10 under H₂ for 4.5h. After purging with nitrogen, the mixture was filtered through a celite pad and the filtrate was concentrated to give the desired title compound **10a**. ¹H NMR (400 MHz, acetone-d₆) δ 7.57-7.47 (m, 5H), 4.95 (m, 1H), 4.73 (d, 1H, OH), 4.06 (m, 1H), 3.65 (m, 1H), 3.60 (m, 1H), 2.90 (m, 1H), 2.25 (t, 2H), 2.25-2.10 (m, 3H), 1.90 (m, 1H), 1.70-1.20 (m, 12H), 1.20 (d, 6H).

15

EXAMPLE 11

7-[(2*S*)-2-[(3*R*)-4,4-difluoro-3-hydroxy-4-phenylbutyl]-5-oxopyrrolidin-1-yl]heptanoic acid (**11a**)

A mixture of compound **10a** (30 mg) and LiOH (0.8 mL, 1M) in
20 MeOH (2.5 mL) was stirred at room temperature for three days and neutralized with 1N HCl. The MeOH was removed in vacuo and the residue extracted with ethyl acetate (3x). The organic layer was washed with water and brine, dried over Na₂SO₄ and filtered. The filtrate was concentrated to give the title compound **11a**. ¹H NMR
25 (400 MHz, acetone-d₆) δ 7.57-7.47 (m, 5H), 4.80 (br s, 1H, OH), 4.06 (m, 1H), 3.68 (m, 1H), 3.52 (m, 1H), 2.91 (m, 1H), 2.28 (t, 2H), 2.25-2.10 (m, 3H), 1.90 (m, 1H), 1.70-1.20 (m, 12H).

30

I. Effects of an EP4 Agonist on Intraocular Pressure (IOP) in Rabbits and Monkeys.

Animals

Drug-naïve, male Dutch Belted rabbits and female cynomolgus
monkeys are used in this study. Animal care and treatment in this investigation are in
compliance with guidelines by the National Institute of Health (NIH) and the
35 Association for Research in Vision and Ophthalmology (ARVO) resolution in the use

of animals for research. All experimental procedures str approved by the Institutional Animal Care and Use Committee of Merck and Company.

Drug Preparation and Administration

- 5 Drug concentrations are expressed in terms of the active ingredient (base). The compounds of this invention are dissolved in physiological saline at 0.01, 0.001, 0.0001 % for rabbit study and 0.05, 0.005% for monkey studies. Drug or vehicle aliquots (25 ul) are administered topically unilaterally or bilaterally. In unilateral applications, the contralateral eyes receive an equal volume of saline.
- 10 Proparacaine (0.5%) is applied to the cornea prior to tonometry to minimize discomfort. Intraocular pressure (IOP) is recorded using a pneumatic tonometer (Alcon Applanation Pneumatonomograph) or equivalent.

Statistical Analysis

- 15 The results are expressed as the changes in IOP from the basal level measured just prior to administration of drug or vehicle and represent the mean, plus or minus standard deviation. Statistical comparisons are made using the Student's t-test for non-paired data between responses of drug-treated and vehicle-treated animals and for paired data between ipsilateral and contralateral eyes at comparable time
- 20 intervals. The significance of the date is also determined as the difference from the "t-0" value using Dunnett's "t" test. Asterisks represent a significance level of $p < 0.05$.

A. Intraocular Pressure Measurement in Rabbits

- 25 Male Dutch Belted rabbits weighing 2.5-4.0 kg are maintained on a 12-hour light/dark cycle and rabbit chow. All experiments are performed at the same time of day to minimize variability related to diurnal rhythm. IOP is measured before treatment then the compounds of this invention or vehicle are instilled (one drop of 25 ul) into one or both eyes and IOP is measured at 30, 60, 120, 180, 240, 300, and 360 minutes after instillation. In some cases, equal number of animals treated bilaterally
- 30 with vehicle only are evaluated and compared to drug treated animals as parallel controls.

B. Intraocular Pressure Measurements in Monkeys.

- 35 Unilateral ocular hypertension of the right eye is induced in female cynomolgus monkeys weighing between 2 and 3 kg by photocoagulation of the

trabecular meshwork with an argon laser system (Coherent NOVUS 2000, Palo Alto, USA) using the method of Lee et al. (1985). The prolonged increase in intraocular pressure (IOP) results in changes to the optic nerve head that are similar to those found in glaucoma patients.


5 For IOP measurements, the monkeys are kept in a sitting position in restraint chairs for the duration of the experiment. Animals are lightly anesthetized by the intramuscular injection of ketamine hydrochloride (3-5 mg/kg) approximately five minutes before each IOP measurement and one drop of 0.5% proparacaine was instilled prior to recording IOP. IOP is measured using a pneumatic tonometer (Alcon
10 Applanation Tonometer) or a Digilab pneumatonometer (Bio-Rad Ophthalmic Division, Cambridge, MA, USA).

IOP is measured before treatment and generally at 30, 60, 124, 180, 300, and 360 minutes after treatment. Baseline values are also obtained at these time points generally two or three days prior to treatment. Treatment consists of instilling
15 one drop of 25 μ l of the compounds of this invention (0.05 and 0.005 %) or vehicle (saline). At least one-week washout period is employed before testing on the same animal. The normotensive (contralateral to the hypertensive) eye is treated in an exactly similar manner to the hypertensive eye. IOP measurements for both eyes are compared to the corresponding baseline values at the same time point. Results are
20 expressed as mean plus-or-minus standard deviation in mm Hg. The activity range of the compounds of this invention for ocular use is between 0.01 and 100,000 nM

By way of example, Compound 7 was a high affinity ligand at the EP4 receptor with a binding affinity in the range of 0.2-2 nM. It was also highly selective against other prostanoid receptors with binding affinities greater than 2 μ M. In a
25 PanLab screening against more than 80 receptors and enzymes, this compound displayed no significant activities at concentrations greater than 10 μ M. The compound was a full agonist at the EP4 receptor with an EC₅₀ of 0.2 -10 nM in a number of cell based functional assays using standard methods for determining EP4 functional agonism. Compound 7 also had good oral bioavailability and terminal
30 elimination half life in rats which are not known for prostaglandin analogs. This unique property allows for convenient oral dosing regimes for study EP4 agonism in in vivo models.

WHAT IS CLAIMED IS:

1. A method for treating ocular hypertension or glaucoma comprising administration to a patient in need of such treatment a therapeutically effective amount of a compound of formula I:





FORMULA I

10

or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug or mixture thereof, wherein,

X is $(CH_2)_n$, O or S;

15

Y represents $(C(R^b)_2)_n$, triple bond,  or  ;

20

R_1 represents hydroxy, CN, CHO, $NHSO_2R_6$, $CONHSO_2R_6$, $CON(R_6)_2$ hydroxymethylketone, $(CH_2)_pCO_2R_6$, $(CH_2)_nSO_3R_6$, C₁₋₄ alkoxy, or $(CH_2)_nC_5-10$ heterocyclyl, said heterocyclyl unsubstituted or substituted with 1 to 3 groups of R_a and optionally containing an acidic hydroxyl group, with the proviso that when X is a bond R_1 is not $(CH_2)_pCO_2R_6$, C₁₋₄ alkoxy, $-(CH_2)_nNR_6R_7$, CHO, $NHSO_2R_6$, $CONHSO_2R_6$, $CON(R_6)_2$, or hydroxymethylketone;

25

R_2 and R_3 independently represents hydrogen, or C₁₋₄ alkyl;

R₆ and R₇ independently represents hydrogen, or C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, (CH₂)_pC₆₋₁₀aryl, (CH₂)_pC₅₋₁₀heterocyclyl, CR²R³OC(O)OC₃₋₁₀cycloalkyl or CR²R³OC(O)O C₁₋₁₀alkyl;

- 5 Ar₂ independently represent (CH₂)_mC₆₋₁₀aryl, (CH₂)_mC₅₋₁₀heteroaryl, (CH₂)_mC₃₋₁₀ heterocycloalkyl, (CH₂)_mC₃₋₈ cycloalkyl said cycloalkyl, heterocycloalkyl, aryl or heteroaryl unsubstituted or substituted with 1-3 groups of R_a;

10 R_a represents C₁₋₆ alkoxy, C₁₋₆ alkyl, CF₃, nitro, amino, cyano, C₁₋₆ alkylamino, or halogen;

R^b independently represents H, halogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or

15 \equiv represents a double or single bond;

p represents 1-3;



n represents 0-4; and

20 m represents 0-8.

2. The method according to claim 1 wherein R₁ is CN, (CH₂)_nC₅₋₁₀heterocyclyl, (CH₂)_pCO₂R₆ or (CH₂)_nSO₃R₆, said heterocyclyl
25 unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described.

3. The method according to claim 2 wherein X and Y are (CH₂)_n.

- 30 4. The method according to claim 1 wherein Y is a double bond as



described by  or  and all other variables are as originally described.

5. The method according to claim 1 wherein R_1 is $(CH_2)_n C_5$ -
10 heterocyclyl, said heterocyclyl unsubstituted or substituted with 1 to 3 groups of R_a , X is $(CH_2)_n$, and Y is $(CH_2)_n$ or $C(halo)_2$.

6. The method according to claim 1 wherein R_1 is
5 $(CH_2)_p CO_2 R_6$, X is $(CH_2)_n$, and Y is $(CH_2)_n$.

7. The method according to claim 1 wherein Ar_2 is $(CH_2)_m C_6$ -
10 aryl, said aryl unsubstituted or substituted with 1 to 3 groups of R_a and all other variables are as originally described.

8. The method according to claim 1 wherein R_1 is a tetrazole
10 unsubstituted or substituted with an R_a group X is $(CH_2)_n$, and Y is $(CH_2)_n$, $C(halo)_2$

or a double bond as described by  or .

9. The method according to claim 1 wherein Ar_2 is a phenyl
unsubstituted or substituted with 1 to 3 groups of R_a , R_1 is tetrazolyl, said tetrazolyl
unsubstituted or substituted with a R_a group and phenyl is unsubstituted or substituted
15 with 1-3 groups of R_a , and all other variables are as originally described.

10. The method according to claim wherein the compound is:
(5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-{4-[(1-methyl-1*H*-
tetrazol-5-yl)]butyl}pyrrolidin-2-one,
4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-
20 yl}butyl cyanate,
3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-
yl}butyl]propanoic acid,
[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-
yl}butyl]methanesulfonic acid,
25 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-{4-[1*H*-tetrazol-5-
ylmethyl]butyl}pyrrolidin-2-one,
[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-
yl}butyl]acetic acid,
(5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-{4-[(1-methyl-1*H*-
30 tetrazol-5-yl)thio]butyl}pyrrolidin-2-one,
(5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(1*H*-tetrazol-5-
ylthio)butyl]pyrrolidin-2-one,

- 3-[4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butylthio]propanoic acid,
 [4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butylthio]methanesulfonic acid,
- 5 (5*R*)-5-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one,
 [4-{(2*R*)-2-[(1*E*)-3-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl}butylthio]acetic acid,
 (5*R*)-5[(1*E*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-1-[6-(1*H*-tetrazol-5-
- 10 yl)hexyl]pyrrolidin-2-one ,
 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 15 7-[(2*S*)-2-[(3*R*)-4,4-difluoro-3-hydroxy-4-phenylbutyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 (5*Z*)-7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]hept-5-enoic acid,
 isopropyl (5*Z*)-7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-
- 20 oxopyrrolidin-1-yl]hept-5-enoate,
 7-[(2*R*)-2-[(1*E*,3*R*)-4-(3-chlorophenyl)-4,4-difluoro-3-hydroxybut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4-(3-chlorophenyl)-4,4-difluoro-3-hydroxybut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 25 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-[3-(trifluoromethyl)phenyl]but-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-[3-(trifluoromethyl)phenyl]but-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
 cyclopentyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-
- 30 oxopyrrolidin-1-yl]heptanoate,
 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-3-methyl-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoic acid,
 isopropyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-3-methyl-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,

- isobutyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
cyclohexyl 7-[(2*R*)-2-[(1*E*,3*R*)-4,4-difluoro-3-hydroxy-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]heptanoate,
- 5 (5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)]butyl]pyrrolidin-2-one,
4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl cyanate,
3-[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]propanoic acid,
- 10 [4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]methanesulfonic acid,
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[1*H*-tetrazol-5-ylmethyl]butyl]pyrrolidin-2-one,
- 15 [4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]acetic acid,
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-[(1-methyl-1*H*-tetrazol-5-yl)thio]butyl]pyrrolidin-2-one,
(5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(1*H*-tetrazol-5-ylthio)butyl]pyrrolidin-2-one,
- 20 3-[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]propanoic acid,
[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]methanesulfonic acid,
- 25 (5*R*)-5-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-1-[4-(methylsulfonyl)butyl]-pyrrolidin-2-one,
[4-[(2*R*)-2-[(1*E*)-(3*R*)-hydroxy-4,4-difluoro-4-phenylbut-1-enyl]-5-oxopyrrolidin-1-yl]butyl]thio]acetic acid, or
(5*R*)-5-[(1*E*)-4,4-difluoro-(3*R*)-hydroxy-4-phenylbut-1-enyl]-1-[6-(1*H*-tetrazol-5-yl)hexyl]pyrrolidin-2-one, a pharmaceutically acceptable salt, enantiomer,
- 30 diastereomer, prodrug, or mixture thereof.

11. A method according to claim 1, which is administered in a topical formulation as a solution or suspension.

12. A method according to claim 1 wherein a second active ingredient belonging to the group consisting of: β -adrenergic blocking agent, parasympatho-mimetic agent, sympathomimetic agent, carbonic anhydrase inhibitor, Maxi-K channel blocker, and a prostaglandin, hypotensive lipid, neuroprotectant, and
5 5-HT₂ receptor agonist is added to the topical formulation.

13. A method according to claim 12 wherein the β -adrenergic blocking agent is timolol, betaxolol, levobetaxolol, carteolol, or levobunolol; the parasympathomimetic agent is pilocarpine; the sympathomimetic agent is
10 epinephrine, brimonidine, iopidine, clonidine, or para-aminoclonidine, the carbonic anhydrase inhibitor is dorzolamide, acetazolamide, metazolamide or brinzolamide; the prostaglandin is latanoprost, travaprost, unoprostone, rescula, or S1033, the hypotensive lipid is lumigan, the neuroprotectant is eliprodil, R-eliprodil or memantine; and the 5-HT₂ receptor agonist is 1-(2-aminopropyl)-3-methyl-1H-
15 imdazol-6-ol fumarate or 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine.


14. A method for treating macular edema, macular degeneration, treating dry eye, increasing retinal and optic nerve head blood velocity, increasing retinal and optic nerve oxygen tension or providing a neuroprotection comprising
20 administration to a patient in need of such treatment a pharmaceutically effective amount of a compound of formula I as recited in claim 1

15. The method according to Claim 14 wherein the compound of formula I is applied as a topical formulation and an active ingredient belonging to the group consisting of β -adrenergic blocking agent, parasympatho-
25 mimetic agent, sympathomimetic agent, carbonic anhydrase inhibitor, Maxi-K channel blocker and a prostaglandin, hypotensive lipid, neuroprotectant, and 5-HT₂ receptor agonist is added to the formulation.

16. A method according to claim 15 wherein the the β -adrenergic blocking agent is timolol, betaxolol, levobetaxolol, carteolol, or levobunolol; the parasympathomimetic agent is pilocarpine; the sympathomimetic agent is epinephrine
30 brimonidine, iopidine, clonidine, or para-aminoclonidine, the carbonic anhydrase inhibitor is dorzolamide, acetazolamide, metazolamide or brinzolamide; the prostaglandin is latanoprost, travaprost, unoprostone, rescula, or S1033, the
35


hypotensive lipid is lumigan, the neuroprotectant is eliprodil, R-eliprodil or memantine; and the 5-HT₂ receptor agonist is 1-(2-aminopropyl)-3-methyl-1H-imidazol-6-ol fumarate or 2-(3-chloro-6-methoxy-indazol-1-yl)-1-methyl-ethylamine.

5 17. A compound of structural formula I:



I

10 or a pharmaceutically acceptable salt, enantiomer, diastereomer, pro drug or mixture thereof, wherein
X is $(CH_2)_n$, O or S;



Ar₂ independently represent (CH₂)_mC₆₋₁₀aryl, (CH₂)_mC₅₋₁₀heteroaryl, (CH₂)_mC₃₋₁₀ heterocycloalkyl, (CH₂)_mC₃₋₈ cycloalkyl said cycloalkyl, heterocycloalkyl, aryl or heteroaryl unsubstituted or substituted with 1-3 groups of R_a;

20 R_a represents C₁₋₆ alkoxy, C₁₋₆ alkyl, CF₃, nitro, amino, cyano, C₁₋₆ alkylamino, or halogen;

R^b independently represents H, halogen, C₁₋₆ alkyl, C₃₋₆ cycloalkyl or

25 --- represents a double or single bond;

n represents 0-4; and

m represents 0-8.

18. The compound according to claim 17 wherein X and Y are $(CH_2)_n$, \equiv represents a double bond; and AR_2 is phenyl.

19. The compound according to claim 18 wherein X is $(CH_2)_n$ and n is 1 and Y is $(CH_2)_n$ and n is 3.

20. Use of a compound of formula I, as defined in any one of claims 1 to 10, or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof, in the manufacture of a medicament for treating hypertension or glaucoma.

21. A pharmaceutical composition for treating hypertension or glaucoma comprising a therapeutically effective amount of a compound of formula I, as defined in any one of claims 1 to 10, or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof, in association with a pharmaceutically acceptable carrier.

22. A composition according to claim 21 in a form for topical administration as a solution or suspension and further comprising a second active ingredient as defined in claim 12 or 13.

23. Use of a compound of formula I, as defined in any one of claims 1 to 10, or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof, in the manufacture of a medicament for treating macular edema, macular degeneration, dry eye, increasing retinal and optic nerve velocity, increasing retinal and optic nerve oxygen tension or providing a neuroprotection.

24. A pharmaceutical composition for treating macular edema, macular degeneration, dry eye, increasing retinal and optic nerve velocity, increasing retinal and optic nerve oxygen tension or providing a neuroprotection comprising a therapeutically effective amount of a compound of formula I, as defined in any one of claims 1 to 10, or a pharmaceutically acceptable salt, enantiomer, diastereomer, prodrug, or mixture thereof, in association with a pharmaceutically acceptable carrier.

25. A composition according to claim 24 in a form for topical administration and further comprising an active ingredient as defined in claim 15 or 16.

26. A compound of claim 17, 18 or 19 for use in medicinal therapy.

INTERNATIONAL SEARCH REPORT

Inter Application No
PCT/CA 03/00838

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P27/06 A61K31/4015 A61K31/41 C07D403/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61P A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DATABASE WPI Section Ch, Week 200247 Derwent Publications Ltd., London, GB; Class B03, AN 2002-443972 XP002252111 -& WO 02 24647 A (ONO PHARM CO LTD), 28 March 2002 (2002-03-28) cited in the application	1-13, 15-22, 26
X	abstract	14, 23-25
Y	WO 02 42268 A (PFIZER PROD INC ;CAMERON KIMBERLY O KEEFE (US); LEFKER BRUCE ALLEN) 30 May 2002 (2002-05-30) cited in the application page 1, line 1; claim 1	17-26

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

22 August 2003

Date of mailing of the international search report

11/09/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Johnson, C

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/CA 03/00838

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00 38667 A (ALCON LAB INC ;KLIMKO PETER G (US); SHARIF NAJAM A (US); GRIFFIN B) 6 July 2000 (2000-07-06) cited in the application page 10, line 6-10 ----	1-13, 20-22
Y	WO 00 38663 A (SHARIF NAJAM A ;ALCON LAB INC (US)) 6 July 2000 (2000-07-06) cited in the application page 6, line 11-13 -----	15,16, 23-25

INTERNATIONAL SEARCH REPORT

Application No.
PCT/CA 03/00838

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1,10,17,20,21,23,24 (all part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,10,17,20,21,23,24 (all part)

Present claims 1, 10, 17, 20, 21, 23, 24 relate to products and methods defined by reference to a desirable characteristic or property, namely prodrugs. The claims cover all products and methods having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products and methods. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/method by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the products of formula I and methods using compounds of formula I, their pharmaceutically acceptable salts, enantiomers, diastereomers and mixtures thereof.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 03/00838

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0224647	A	28-03-2002	AU 9025001 A WO 0224647 A1	02-04-2002 28-03-2002
WO 0242268	A	30-05-2002	AU 1084802 A CA 2429850 A1 WO 0242268 A2 US 2003149086 A1 US 2002065308 A1	03-06-2002 30-05-2002 30-05-2002 07-08-2003 30-05-2002
WO 0038667	A	06-07-2000	AU 2211700 A WO 0038667 A2 US 6545045 B1	31-07-2000 06-07-2000 08-04-2003
WO 0038663	A	06-07-2000	AU 2183900 A WO 0038663 A2 US 6344477 B1	31-07-2000 06-07-2000 05-02-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.